

# Best Available Copy



**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : <b>C12N 15/861, 15/62, 15/34, 5/10, C07K 14/075, A61K 48/00, A61P 9/00</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 00/31285</b></p> <p>(43) International Publication Date: <b>2 June 2000 (02.06.00)</b></p>
<p>(21) International Application Number: <b>PCT/NL99/00717</b></p> <p>(22) International Filing Date: <b>22 November 1999 (22.11.99)</b></p> <p>(30) Priority Data: <b>98203921.6 20 November 1998 (20.11.98) EP</b></p> <p>(71) Applicant: <b>INTROGENE B.V. [NL/NL]; Wassenaarseweg 72, NL-2333 AL Leiden (NL).</b></p> <p>(72) Inventors: <b>HAVENGA, Menzo, Jans, Emco; Stortemelk 57, NL-2401 BT Alphen aan de Rijn (NL). BOUT, Abraham; Coymansstraat 24, NL-2751 AR Moerkapelle (NL). VOGELS, Ronald; Van Rietlaan 4, NL-3461 HW Linschoten (NL).</b></p> <p>(74) Agent: <b>OTTEVANGERS, S., U.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).</b></p>		<p>(81) Designated States: <b>AM, AZ, BA, BB, BG, BR, BY, CA, CN, CR, CU, CZ, DM, GD, GE, GH, GM, HR, HU, ID, IN, IS, KE, KG, KP, KR, KZ, LC, LK, LR, LS, MA, MD, MG, MN, MW, PL, RU, SD, SG, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: <b>GENE DELIVERY VECTORS PROVIDED WITH A TISSUE TROPISM FOR SMOOTH MUSCLE CELLS, AND/OR ENDOTHELIAL CELLS</b></p>		
<p>(57) Abstract</p> <p>The invention provides a nucleic acid delivery vehicle with or having been provided with at least a tissue tropism for smooth muscle cells and/or endothelial cells. In one aspect said nucleic acid delivery vehicle is a virus capsid or a functional part, derivative and/or analogue thereof. Preferably said virus capsid is an adenovirus capsid. Preferably said adenovirus is a subgroup B adenovirus, preferably adenovirus 16. Preferably said tissue tropism is provided by at least a tissue tropism determining part of an adenovirus fiber protein or a functional derivative and/or analogue thereof. The invention further presents methods for the treatment of diseases, preferably cardiovascular diseases.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Gene delivery vectors provided with a tissue tropism for smooth muscle cells, and/or endothelial cells.

#### FIELD OF THE INVENTION

The invention relates to the field of molecular genetics and medicine. In particular the present invention relates to the field of gene therapy, more in particular to  
5 gene therapy using adenoviruses.

#### BACKGROUND OF THE INVENTION

In gene therapy, genetic information is usually delivered to a host cell in order to either correct  
10 (supplement) a genetic deficiency in said cell, or to inhibit an undesired function in said cell, or to eliminate said host cell. Of course the genetic information can also be intended to provide the host cell with a desired function, e.g. to supply a secreted protein to treat other  
15 cells of the host, etc.

Many different methods have been developed to introduce new genetic information into cells. Although many different systems may work on cell lines cultured in vitro, only the  
20 group of viral vector mediated gene delivery methods seems to be able to meet the required efficiency of gene transfer in vivo. Thus for gene therapy purposes most of the attention is directed toward the development of suitable viral vectors. Today, most of the attention for the  
25 development of suitable viral vectors is directed toward those vectors that are based on adenoviruses. These adenovirus vectors can deliver foreign genetic information very efficiently to target cells in vivo. Moreover, obtaining large amounts of adenovirus vectors is for most  
30 types of adenovirus vectors not a problem. Adenovirus vectors are relatively easy to concentrate and purify. Moreover, studies in clinical trials have provided valuable information on the use of these vectors in patients.

There are a lot of reasons for using adenovirus vectors for the delivery of nucleic acid to target cells in gene therapy protocols. However, some characteristics of the current vectors limit their use in specific applications. For instance endothelial cells and smooth muscle cells are not easily transduced by the current generation of adenovirus vectors. For many gene therapy applications, such as applications in the cardiovascular area, preferably these types of cells should be genetically modified. On the other hand, in some applications, even the very good in vivo delivery capacity of adenovirus vectors is not sufficient and higher transfer efficiencies are required. This is the case, for instance, when most cells of a target tissue need to be transduced.

The present invention was made in the course of the manipulation of adenovirus vectors. In the following section therefore a brief introduction to adenoviruses is given.

20

#### **Adenoviruses**

Adenoviruses contain a linear double-stranded DNA molecule of approximately 36000 base pairs. It contains identical Inverted Terminal Repeats (ITR) of approximately 90-140 base pairs with the exact length depending on the serotype. The viral origins of replication are within the ITRs exactly at the genome ends. The transcription units are divided in early and late regions. Shortly after infection the E1A and E1B proteins are expressed and function in transactivation of cellular and adenoviral genes. The early regions E2A and E2B encode proteins (DNA binding protein, pre-terminal protein and polymerase) required for the replication of the adenoviral genome (reviewed in van der Vliet, 1995). The early region E4 encodes several proteins with pleiotropic functions e.g. transactivation of the E2 early promoter, facilitating transport and accumulation of viral mRNAs in the late phase of infection and increasing

nuclear stability of major late pre-mRNAs (reviewed in Leppard, 1997). The early region 3 encodes proteins that are involved in modulation of the immune response of the host (Wold et al, 1995). The late region is transcribed from one  
5 single promoter (major late promoter) and is activated at the onset of DNA replication. Complex splicing and poly-adenylation mechanisms give rise to more than 12 RNA species coding for core proteins, capsid proteins (penton, hexon, fiber and associated proteins), viral protease and proteins  
10 necessary for the assembly of the capsid and shut-down of host protein translation (Imperiale, M.J., Akusjnarvi, G. and Leppard, K.N. (1995) Post-transcriptional control of adenovirus gene expression. In: The molecular repertoire of adenoviruses I. P139-171. W. Doerfler and P. Bohm (eds),  
15 Springer-Verlag Berlin Heidelberg).

#### *Interaction between virus and host cell*

The interaction of the virus with the host cell has mainly been investigated with the serotype C viruses Ad2 and  
20 Ad5. Binding occurs via interaction of the knob region of the protruding fiber with a cellular receptor. The receptor for Ad2 and Ad5 and probably more adenoviruses is known as the 'Coxsackievirus and Adenovirus Receptor' or CAR protein (Bergelson et al, 1997). Internalization is mediated through  
25 interaction of the RGD sequence present in the penton base with cellular integrins (Wickham et al, 1993). This may not be true for all serotypes, for example serotype 40 and 41 do not contain a RGD sequence in their penton base sequence (Kidd et al, 1993).

30

#### *The fiber protein*

The initial step for successful infection is binding of adenovirus to its target cell, a process mediated through fiber protein. The fiber protein has a trimeric structure  
35 (Stouten et al, 1992) with different lengths depending on

the virus serotype (Signas et al, 1985; Kidd et al, 1993). Different serotypes have polypeptides with structurally similar N and C termini, but different middle stem regions. The first 30 amino acids at the N terminus are involved in anchoring of the fiber to the penton base (Chroboczek et al, 1995), especially the conserved FNPVYP region in the tail (Arnberg et al, 1997). The C-terminus, or knob, is responsible for initial interaction with the cellular adenovirus receptor. After this initial binding secondary binding between the capsid penton base and cell-surface integrins leads to internalization of viral particles in coated pits and endocytosis (Morgan et al, 1969; Svensson and Persson, 1984; Varga et al, 1991; Greber et al, 1993; Wickham et al, 1993). Integrins are  $\alpha\beta$ -heterodimers of which at least 14  $\alpha$ -subunits and 8  $\beta$ -subunits have been identified (Hynes, 1992). The array of integrins expressed in cells is complex and will vary between cell types and cellular environment. Although the knob contains some conserved regions, between serotypes, knob proteins show a high degree of variability, indicating that different adenovirus receptors exist.

#### *Adenoviral serotypes*

At present, six different subgroups of human adenoviruses have been proposed which in total encompass approximately 50 distinct adenovirus serotypes. Besides these human adenoviruses, many animal adenoviruses have been identified (see e.g. Ishibashi and Yasue, 1984). A serotype is defined on the basis of its immunological distinctiveness as determined by quantitative neutralization with animal antiserum (horse, rabbit). If neutralization shows a certain degree of cross-reaction between two viruses, distinctiveness of serotype is assumed if A) the hemagglutinins are unrelated, as shown by lack of cross-reaction on hemagglutination-inhibition, or B) substantial biophysical/biochemical differences in DNA exist (Francki et

al, 1991). The serotypes identified last (42-49) were isolated for the first time from HIV infected patients (Hierholzer et al, 1988; Schnurr et al, 1993). For reasons not well understood, most of such immuno-compromised patients shed adenoviruses that were never isolated from immuno-competent individuals (Hierholzer et al, 1988, 1992; Khoo et al, 1995).

Besides differences towards the sensitivity against neutralizing antibodies of different adenovirus serotypes, adenoviruses in subgroup C such as Ad2 and Ad5 bind to different receptors as compared to adenoviruses from subgroup B such as Ad3 and Ad7 (Defer et al, 1990; Gall et al, 1996). Likewise, it was demonstrated that receptor specificity could be altered by exchanging the Ad3 knob protein with the Ad 5 knob protein, and vice versa (Krasnykh et al, 1996; Stevenson et al, 1995, 1997). Serotypes 2, 4, 5 and 7 all have a natural affiliation towards lung epithelia and other respiratory tissues. In contrast, serotypes 40 and 41 have a natural affiliation towards the gastrointestinal tract. These serotypes differ in at least capsid proteins (penton-base, hexon), proteins responsible for cell binding (fiber protein), and proteins involved in adenovirus replication. It is unknown to what extent the capsid proteins determine the differences in tropism found between the serotypes. It may very well be that post-infection mechanisms determine cell type specificity of adenoviruses. It has been shown that adenoviruses from serotypes A (Ad12 and Ad31), C (Ad2 and Ad5), D (Ad9 and Ad15), E (Ad4) and F (Ad41) all are able to bind labeled, soluble CAR (sCAR) protein when immobilized on nitrocellulose. Furthermore, binding of adenoviruses from these serotypes to Ramos cells, that express high levels of CAR but lack integrins (Roelvink et al, 1996), could be efficiently blocked by addition of sCAR to viruses prior to infection (Roelvink et al, 1998). However, the fact that (at least some) members of these

subgroups are able to bind CAR does not exclude that these viruses have different infection efficiencies in various cell types. For example subgroup D serotypes have relatively short fiber shafts compared to subgroup A and C viruses. It has been postulated that the tropism of subgroup D viruses is to a large extent determined by the penton base binding to integrins (Roelvink et al, 1996; Roelvink et al, 1998). Another example is provided by Zabner et al, 1998 who have tested 14 different serotypes on infection of human ciliated airway epithelia (CAE) and found that serotype 17 (subgroup D) was bound and internalized more efficiently than all other viruses, including other members of subgroup D. Similar experiments using serotypes from subgroup A-F in primary fetal rat cells showed that adenoviruses from subgroup A and B were inefficient whereas viruses from subgroup D were most efficient (Law et al, 1998). Also in this case viruses within one subgroup displayed different efficiencies. The importance of fiber binding for the improved infection of Ad17 in CAE was shown by Armentano et al (WO 98/22609) who made a recombinant LacZ Ad2 virus with a fiber gene from Ad17 and showed that the chimaeric virus infected CAE more efficiently than LacZ Ad2 viruses with Ad2 fibers.

Thus despite their shared ability to bind CAR, differences in the length of the fiber, knob sequence and other capsid proteins e.g. penton base of the different serotypes may determine the efficiency by which an adenovirus infects a certain target cell. Of interest in this respect is the ability of Ad5 and Ad2 fibers but not of Ad3 fibers to bind to fibronectin III and MHC class 1  $\alpha 2$  derived peptides. This suggests that adenoviruses are able to use cellular receptors other than CAR (Hong et al, 1997). Serotypes 40 and 41 (subgroup F) are known to carry two fiber proteins differing in the length of the shaft. The long shafted 41L fiber is shown to bind CAR whereas the

short shafted 41S is not capable of binding CAR (Roelvink et al, 1998). The receptor for the short fiber is not known.

#### *Adenoviral gene delivery vectors*

5        Most adenoviral gene delivery vectors currently used in gene therapy are derived from the serotype C adenoviruses Ad2 or Ad5. The vectors have a deletion in the E1 region, where novel genetic information can be introduced. The E1 deletion renders the recombinant virus replication  
10 defective. It has been demonstrated extensively that recombinant adenovirus, in particular serotype 5 is suitable for efficient transfer of genes in vivo to the liver, the airway epithelium and solid tumors in animal models and human xenografts in immuno-deficient mice (Bout 1996, 1997;  
15 Blaese et al, 1995).

Gene transfer vectors derived from adenoviruses (adenoviral vectors) have a number of features that make them particularly useful for gene transfer:

- 20 1) the biology of the adenoviruses is well characterized,
- 2) the adenovirus is not associated with severe human pathology,
- 3) the virus is extremely efficient in introducing its DNA into the host cell,
- 25 4) the virus can infect a wide variety of cells and has a broad host-range,
- 5) the virus can be produced at high titers in large quantities,
- 6) and the virus can be rendered replication defective by  
30 deletion of the early-region 1 (E1) of the viral genome (Brody and Crystal, 1994).

However, there is still a number of drawbacks associated with the use of adenoviral vectors:

1) Adenoviruses, especially the well investigated serotypes Ad2 and Ad5 usually elicit an immune response by the host into which they are introduced,

2) it is currently not feasible to target the virus to  
5 certain cells and tissues,

3) the replication and other functions of the adenovirus are not always very well suited for the cells, which are to be provided with the additional genetic material,

4) the serotypes Ad2 or Ad5, are not ideally suited for  
10 delivering additional genetic material to organs other than the liver. The liver can be particularly well transduced with vectors derived from Ad2 or Ad5. Delivery of such vectors via the bloodstream leads to a significant deliver of the vectors to the cells of the liver. In therapies were  
15 other cell types then liver cells need to be transduced some means of liver exclusion must be applied to prevent uptake of the vector by these cells. Current methods rely on the physical separation of the vector from the liver cells, most of these methods rely on localizing the vector and/or the  
20 target organ via surgery, balloon angioplasty or direct injection into an organ via for instance needles. Liver exclusion is also being practiced through delivery of the vector to compartments in the body that are essentially isolated from the bloodstream thereby preventing transport  
25 of the vector to the liver. Although these methods mostly succeed in avoiding gross delivery of the vector to the liver, most of the methods are crude and still have considerable leakage and/or have poor target tissue penetration characteristics. In some cases inadvertent  
30 delivery of the vector to liver cells can be toxic to the patient. For instance, delivery of a herpes simplex virus (HSV) thymidine kinase (TK) gene for the subsequent killing of dividing cancer cells through administration of gancyclovir is quite dangerous when also a significant  
35 amount of liver cells are transduced by the vector. Significant delivery and subsequent expression of the HSV-TK gene to liver cells is associated with severe toxicity. Thus

there is a discrete need for an inherently safe vector provided with the property of a reduced transduction efficiency of liver cells.

## 5 BRIEF DESCRIPTION OF DRAWINGS

Table I: Oligonucleotides and degenerate oligonucleotides used for the amplification of DNA encoding fiber proteins derived from alternative adenovirus serotypes. (Bold letters represent NdeI restriction site (A-E), NsiI restriction site (1-6, 8), or PacI restriction site, (7)).

Table II: Biodistribution of chimeric adenovirus upon intravenous tail vein injection. Values represent luciferase activity/  $\mu\text{g}$  of total protein. All values below 200 Relative light units/  $\mu\text{g}$  protein are considered background. ND = not determined

Table III: Expression of CAR and integrins on the cell surface of endothelial cells and smooth muscle cells. 70%: Cells harvested for FACS analysis at a cell density of 70% confluency. 100%: Cells harvested for FACS analysis at a cell density of 100% confluency. PER.C6 cells were taken as a control for antibody staining. Values represent percentages of cells that express CAR or either one of the integrins at levels above background. As background control, HUVECs or HUVsmc were incubated only with the secondary, rat-anti-mouse IgG1-PE labeled antibody.

Table IV: Determination of transgene expression (luciferase activity) per  $\mu\text{g}$  of total cellular protein after infection of A549 cells.

30

Figure 1: Schematic drawing of the pBr/Ad.Bam-rITR construct.

Figure 2: Schematic drawing of the strategy used to delete the fiber gene from the pBr/Ad.Bam-rITR construct.

Figure 3: Schematic drawing of construct pBr/Ad.BamRAfib.

35 Figure 4: Sequences of the chimaeric fibers Ad5/12, Ad5/16, Ad5/28, and Ad5/40-L.

Figure 5: Schematic drawing of the construct pClipsal-Luc.

Figure 6: Schematic drawing of the method to generate chimaeric adenoviruses using three overlapping fragments. Early (E) and late regions (L) are indicated. L5 is the fiber coding sequence.

Figure 7: A) Infection of HUVEC cells using different amounts of virus particles per cell and different fiber chimeric adenoviruses. Virus concentration: 10000 vp/ cell (= white bar), 5000 vp/ cell (= grey bar), 2500 vp/ cell (= Black bar) 1000 vp/ cell (light grey bar, 250 and 50 vp/ cell no detectable luciferase activity above background. Luciferase activity is expressed in relative light units (RLU) per microgram cellular protein. B) Infection of HUVEC cells using different concentrations of cells (22500, 45000, 90000, or 135000 cells seeded per well) and either adenovirus serotype 5 (black bar) or the fiber 16 chimeric adenovirus (white bar). Luciferase activity is expressed in relative light units (RLU) per microgram cellular protein. C) Flow cytometric analysis on Human aorta EC transduced with 500 (Black bar) or 5000 (grey bar) virus particles per cell of Ad5 or the fiber 16 chimeric virus (Fib16). Non-infected cells were used to set the background at 1% and a median fluorescence of 5.4. The maximum shift in the median fluorescence that can be observed on a flow cytometer is 9999. This latter indicates that at 5000 vp/ cell both Ad5 and Fib16 are outside the sensitivity scale of the flow cytometer.

Figure 8: A) Infection of HUVsmc cells using different amounts of virus particles per cell and different fiber mutant Ad5 based adenoviruses. Virus concentration: 5000 vp/ cell (= white bar), 2500 vp/ cell (= grey bar), 1250 vp/ cell (= dark grey bar), 250 vp/ cell (= black bar), or 50 vp/ cell (light grey bar). Luciferase activity is expressed as relative light units (RLU) per microgram cellular protein. B) Infection of HUVsmc cells using different concentrations of cells (10000, 20000, 40000, 60000, or 80000 cells per well) and either adenovirus

serotype 5 (white bars) or the fiber 16 chimeric adenovirus (black bars). A plateau is observed after infection with chimeric fiber 16 adenovirus due to the fact that transgene expression is higher than the sensitivity range of the bioluminimeter used. C) Human umbilical vein SMC transduced with 500 VP/ cell (black bar) or 5000 VP/ cell (grey bar) using either Ad5 or the fiber 16 mutant (Fib16). Non-transduced cells were used to set a background median fluorescence of approximately 1. Shown is the median fluorescence of GFP expression as measured by flow cytometry. D) HUVsmc were infected with 312 (light grey bar), 625 (grey bar), 1250 (black bar), 2500 (dark grey bar), 5000 (light grey bar), or 10000 (white bar) virus particles per cell of either the fiber 11, 16, 35, or 51 chimeric virus. Luciferase transgene expression expressed as relative light units (RLU) per microgram protein was measured 48 hours after virus exposure. E) Macroscopic photographs of LacZ staining on saphenous samples. Nuclear targeted LacZ (ntLacZ) yields a deep blue color which appears black or dark grey in non-color prints. F) Macroscopic photographs of LacZ staining on pericard samples. Nuclear targeted LacZ (ntLacZ) gives a deep blue color which appears black in non-color prints G) Macroscopic photographs of LacZ staining on right coronary artery samples. Nuclear targeted LacZ (ntLacZ) gives a deep blue color which appears black in non-color prints H) LacZ staining on Left artery descending (LAD) samples. Nuclear targeted LacZ (ntLacZ) gives a deep blue color which appears black in non-color prints

Figure 9: Sequences including the gene encoding adenovirus 16 fiber protein as published in Genbank and sequences including a gene encoding a fiber from an adenovirus 16 variant as isolated in the present invention, wherein the sequences of the fiber protein are from the NdeI-site.

Figure 9A nucleotide sequence comparison. Figure 9B amino-acid comparison.

## SUMMARY OF THE INVENTION

The present invention provides gene therapy methods, compounds and medicines. The present invention is particularly useful in gene therapy applications were  
5 endothelial cells and /or smooth muscle cells form the target cell type. The present invention relates to gene delivery vehicles provided with a tissue tropism for at least endothelial cells and /or smooth muscle cells. The present invention further relates to gene delivery vehicles having  
10 been deprived of a tissue tropism for liver cells.

## DETAILED DESCRIPTION OF THE INVENTION.

It is an object of the current invention to provide materials and methods to overcome the limitations of  
15 adenoviral vectors mentioned above. In a broad sense, the invention provides new adenoviruses, derived in whole or in part from serotypes different from Ad5. Specific genes of serotypes with preferred characteristics may be combined in a chimaeric vector to give rise to a vector that is better  
20 suited for specific applications. Preferred characteristics include, but are not limited to, improved infection of a specific target cell, reduced infection of non-target cells, improved stability of the virus, reduced uptake in antigen presenting cells (APC), or increased uptake in APC, reduced  
25 toxicity to target cells, reduced neutralization in humans or animals, reduced or increased CTL response in humans or animals, better and/or prolonged transgene expression, increased penetration capacity in tissues, improved yields in packaging cell lines, etc.

30

One aspect of the present invention facilitates the combination of the low immunogenicity of some adenoviruses with the characteristics of other adenoviruses that allow efficient gene therapy. Such characteristics may be a high

specificity for certain host cells, a good replication machinery for certain cells, a high rate of infection in certain host cells, low infection efficiency in non-target cells, high or low efficiency of APC infection, etc.

- 5 The invention thus may provide chimaeric adenoviruses having the useful properties of at least two adenoviruses of different serotypes.

Typically, two or more requirements from the above non-exhaustive list are required to obtain an adenovirus capable  
10 of efficiently transferring genetic material to a host cell. Therefore the present invention provides adenovirus derived vectors which can be used as cassettes to insert different adenoviral genes from different adenoviral serotypes at the required sites. This way one can obtain a vector capable of  
15 producing a chimaeric adenovirus, whereby of course also a gene of interest can be inserted (for instance at the site of E1 of the original adenovirus). In this manner the chimaeric adenovirus to be produced can be adapted to the requirements and needs of certain hosts in need of gene  
20 therapy for certain disorders. To enable this virus production, a packaging cell will generally be needed in order to produce sufficient amount of safe chimaeric adenoviruses.

In one of its aspects the present invention provides  
25 adenoviral vectors comprising at least a fragment of a fiber protein of an adenovirus from subgroup B. Said fiber protein may be the native fiber protein of the adenoviral vector or may be derived from a serotype different from the serotype the adenoviral vector is based on. In the latter case the  
30 adenoviral vector according to the invention is a chimaeric adenovirus displaying at least a fragment of the fiber protein derived from subgroup B adenoviruses that fragment comprising at least the receptor binding sequence. Typically such a virus will be produced using a vector (typically a  
35 plasmid, a cosmid or baculovirus vector). Such vectors are also subject of the present invention. A preferred vector is a vector that can be used to make a chimaeric recombinant

virus specifically adapted to the host to be treated and the disorder to be treated.

The present invention also provides a chimaeric adenovirus based on adenovirus type 5 but having at least a fragment of  
5 the fiber sequence from adenovirus type 16, whereby the fragment of the fiber of Ad16 comprises the fragment of the fiber protein that is involved in binding a host cell.

The present invention also provides chimaeric adenoviral vectors that show improved infection as compared to  
10 adenoviruses from other subgroups in specific host cells for example, but not limited to, endothelial cells and smooth muscle cells of human or animal origin. An important feature of the present invention is the means to produce the chimaeric virus. Typically, one does not want an adenovirus  
15 batch to be administered to the host cell, which contains replication competent adenovirus. In general therefore it is desired to omit a number of genes (but at least one) from the adenoviral genome on the vector encoding the chimaeric virus and to supply these genes in the genome of the cell in  
20 which the vector is brought to produce chimaeric adenovirus. Such a cell is usually called a packaging cell. The invention thus also provides a packaging cell for producing a chimaeric adenovirus according to the invention, comprising in trans all elements necessary for adenovirus  
25 production not present on the adenoviral vector according to the invention. Typically vector and packaging cell have to be adapted to one another in that they have all the necessary elements, but that they do not have overlapping elements which lead to replication competent virus by  
30 recombination. Thus the invention also provides a kit of parts comprising a packaging cell according to the invention and a recombinant vector according the invention whereby there is essentially no sequence overlap leading to recombination resulting in the production of replication  
35 competent adenovirus between said cell and said vector. For certain applications for example when the therapy is aimed at eradication of tumor cells, the adenoviral vector

according to the invention may be replication competent or capable of replicating under certain conditions for example in specific cell types like tumor cells or tumor endothelial cells.

- 5 It is within the scope of the invention to insert more genes, or a functional part of these genes from the same or other serotypes into the adenoviral vector replacing the corresponding native sequences. Thus for example replacement of (a functional part of the) fiber sequences with
- 10 corresponding sequences of other serotypes may be combined with for example replacements of (a functional part of) other capsid genes like penton base or hexon with corresponding sequences of said serotype or of other distinct serotypes. Persons skilled in the art understand
- 15 that other combinations not limited to the said genes are possible and are within the scope of the invention. The chimaeric adenoviral vector according to the invention may originate from at least two different serotypes. This may provide the vector with preferred characteristics such as
- 20 improved infection of target cells and/or less infection of non-target cells, improved stability of the virus, reduced immunogenicity in humans or animals (e.g. reduced uptake in APC, reduced neutralization in the host and/or reduced cytotoxic T-lymphocyte (CTL) response), increased
- 25 penetration of tissue, better longevity of transgene expression, etc. In this aspect it is preferred to use capsid genes e.g. penton and/or hexon genes from less immunogenic serotypes as defined by the absence or the presence of low amounts of neutralizing antibodies in the
- 30 vast majority of hosts. It is also preferred to use fiber and/or penton sequences from serotypes that show improved binding and internalization in the target cells. Furthermore it is preferred to delete from the viral vector those genes which lead to expression of adenoviral genes in the target
- 35 cells. In this aspect a vector deleted of all adenoviral genes is also preferred. Furthermore it is preferred that

the promoter driving the gene of interest to be expressed in the target cells is a cell type specific promoter.

In order to be able to precisely adapt the viral vector and provide the chimaeric virus with the desired properties at will, it is preferred that a library of adenoviral genes is provided whereby the genes to be exchanged are located on plasmid- or cosmid-based adenoviral constructs whereby the genes or the sequences to be exchanged are flanked by restriction sites. The preferred genes or sequences can be selected from the library and inserted in the adenoviral constructs that are used to generate the viruses. Typically such a method comprises a number of restriction and ligation steps and transfection of a packaging cell. The adenoviral vector can be transfected in one piece, or as two or more overlapping fragments, whereby viruses are generated by homologous recombination. For example the adenoviral vector may be built up from two or more overlapping sequences for insertion or replacements of a gene of interest in for example the E1 region, for insertion or replacements in penton and/or hexon sequences, and for insertions or replacements into fiber sequences. Thus the invention provides a method for producing chimaeric adenoviruses having one or more desired properties like a desired host range and diminished antigenicity, comprising providing one or more vectors according to the invention having the desired insertion sites, inserting into said vectors at least a functional part of a fiber protein derived from an adenovirus serotype having the desired host range and/or inserting a functional part of a capsid protein derived from an adenovirus serotype having relatively low antigenicity and transfecting said vectors in a packaging cell according to the invention and allowing for production of chimaeric viral particles. Of course other combinations of other viral genes originating from different serotypes can also be inserted as disclosed herein before. Chimaeric viruses having only one non-native sequence in addition to an

insertion or replacement of a gene of interest in the E1 region, are also within the scope of the invention.

An immunogenic response to adenovirus that typically occurs is the production of neutralizing antibodies by the host.

- 5 This is typically a reason for selecting a capsid protein like penton, hexon and/or fiber of a less immunogenic serotype.

- Of course it may not be necessary to make chimaeric adenoviruses which have complete proteins from different  
10 serotypes. It is well within the skill of the art to produce chimaeric proteins, for instance in the case of fiber proteins it is very well possible to have the base of one serotype and the shaft and the knob from another serotype. In this manner it becomes possible to have the parts of the  
15 protein responsible for assembly of viral particles originate from one serotype, thereby enhancing the production of intact viral particles. Thus the invention also provides a chimaeric adenovirus according to the invention, wherein the hexon, penton, fiber and/or other  
20 capsid proteins are chimaeric proteins originating from different adenovirus serotypes. Besides generating chimaeric adenoviruses by swapping entire wild type capsid (protein) genes etc. or parts thereof, it is also within the scope of the present invention to insert capsid (protein) genes etc.  
25 carrying non-adenoviral sequences or mutations such as point mutations, deletions, insertions, etc. which can be easily screened for preferred characteristics such as temperature stability, assembly, anchoring, redirected infection, altered immune response etc. Again other chimaeric  
30 combinations can also be produced and are within the scope of the present invention.

- It has been demonstrated in mice and rats that upon in vivo systemic delivery of recombinant adenovirus of common used  
35 serotypes for gene therapy purposes more than 90% of the

virus is trapped in the liver (Herz et al, 1993; Kass-Eisler et al, 1994; Huard et al, 1995). It is also known that human hepatocytes are efficiently transduced by adenovirus serotype 5 vectors (Castell, J.V., Hernandez, D. Gomez-Foix, A.M., Guillen, I, Donato, T. and Gomez-Lechon, M.J. (1997). Adenovirus-mediated gene transfer into human hepatocytes: analysis of the biochemical functionality of transduced cells. Gene Ther. 4(5), p455-464). Thus in vivo gene therapy by systemic delivery of Ad2 or Ad5 based vectors is seriously hampered by the efficient uptake of the viruses in the liver leading to unwanted toxicity and less virus being available for transduction of the target cells. Therefore, alteration of the adenovirus serotype 5 host cell range to be able to target other organs in vivo is a major interest of the invention.

To obtain re-directed infection of recombinant adenovirus serotype 5, several approaches have been or still are under investigation. Wickham et al have altered the RGD (Arg, Gly, Asp) motif in the penton base which is believed to be responsible for the  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$  integrin binding to the penton base. They have replaced this RGD motif by another peptide motif which is specific for the  $\alpha_4\beta_1$  receptor. In this way targeting the adenovirus to a specific target cell could be accomplished (Wickham et al, 1995). Krasnykh et al (1998) have made use of the HI loop available in the knob. This loop is, based on X-ray crystallography, located on the outside of the knob trimeric structure and therefore is thought not to contribute to the intramolecular interactions in the knob. Insertion of a FLAG coding sequence into the HI loop resulted in fiber proteins that were able to trimerise and it was further shown that viruses containing the FLAG sequence in the knob region could be made. Although interactions of the FLAG-containing knob with CAR are not changed, insertion of ligands in the HI loop may lead to retargeting of infection. Although successful introduction

of changes in the adenovirus serotype 5 fiber and penton-base have been reported, the complex structure of knob and the limited knowledge of the precise amino acids interacting with CAR render such targeting approaches laborious and difficult. The use of antibodies binding to CAR and to a specific cellular receptor has also been described (Wickham et al, 1996; Rogers et al, 1997). This approach is however limited by the availability of a specific antibody and by the complexity of the gene therapy product.

10 To overcome the limitations described above we used pre-existing adenovirus fibers, penton base proteins, hexon proteins or other capsid proteins derived from other adenovirus serotypes. By generating chimaeric adenovirus serotype 5 libraries containing structural proteins of

15 alternative adenovirus serotypes, we have developed a technology, which enables rapid screening for a recombinant adenoviral vector with preferred characteristics.

It is an object of the present invention to provide methods for the generation of chimaeric capsids which can be targeted to specific cell types *in vitro* as well as *in vivo*, and thus have an altered tropism for certain cell types. It is a further object of the present invention to provide methods and means by which an adenovirus or an adenovirus capsid can be used as a protein or nucleic acid delivery vehicle to a specific cell type or tissue.

20

25

The generation of chimaeric adenoviruses based on adenovirus serotype 5 with modified late genes is described. For this purpose, three plasmids, which together contain the complete adenovirus serotype 5 genome, were constructed. From one of these plasmids part of the DNA encoding the adenovirus serotype 5 fiber protein was removed and replaced by linker DNA sequences that facilitate easy cloning. This plasmid subsequently served as template for the insertion of DNA encoding fiber protein derived from different adenovirus serotypes. The DNA sequences derived from the different

30

35

serotypes were obtained using the polymerase chain reaction technique in combination with (degenerate) oligonucleotides. At the former E1 location in the genome of adenovirus serotype 5, any gene of interest can be cloned. A single  
5 transfection procedure of the three plasmids together results in the formation of a recombinant chimaeric adenovirus. Alternatively, cloning of the sequences obtained from the library of genes can be such that the chimaeric adenoviral vector is build up from one or two fragments. For  
10 example one construct contains at least the left ITR and sequences necessary for packaging of the virus, an expression cassette for the gene of interest and sequences overlapping with the second construct comprising all sequences necessary for replication and virus formation not  
15 present in the packaging cell as well as the non-native sequences providing the preferred characteristics. This new technology of libraries consisting of chimaeric adenoviruses thus allows for a rapid screening for improved recombinant adenoviral vectors for *in vitro* and *in vivo* gene therapy  
20 purposes.

The use of adenovirus type 5 for *in vivo* gene therapy is limited by the apparent inability to infect certain cell types e.g. human endothelial cells and smooth muscle cells  
25 and the preference of infection of certain organs e.g. liver and spleen. Specifically this has consequences for treatment of cardiovascular diseases like restenosis and pulmonary hypertension. Adenovirus-mediated delivery of human ceNOS (constitutive endothelial nitric oxide synthase) has been  
30 proposed as treatment for restenosis after percutaneous transluminal coronary angioplasty (PTCA). Restenosis is characterized by progressive arterial remodeling, extracellular matrix formation and intimal hyperplasia at the site of angioplasty (Schwartz et al, 1993; Carter et al,  
35 1994; Shi et al, 1996). NO is one of the vasoactive factors shown to be lost after PTCA-induced injury to the

endothelial barrier (Lloyd Jones and Bloch, 1996). Thus restoration of NO levels after balloon-induced injury by means of adenoviral delivery of ceNOS may prevent restenosis (Varenne et al, 1998). Other applications for gene therapy

5 whereby the viruses or chimaeric viruses according to the invention are superior to Ad2 or Ad5 based viruses, given as non-limiting examples, are production of proteins by endothelial cells that are secreted into the blood, treatment of hypertension, preventive treatment of stenosis

10 during vein grafting, angiogenesis, heart failure, renal hypertension and others.

In one embodiment this invention describes adenoviral vectors that are, amongst others, especially suited for gene

15 delivery to endothelial cells and smooth muscle cells important for treatment of cardiovascular disorders. The adenoviral vectors preferably are derived from subgroup B adenoviruses or contain at least a functional part of the fiber protein from an adenovirus from subgroup B comprising

20 at least the cell-binding moiety of the fiber protein.

In a further preferred embodiment the adenoviral vectors are chimaeric vectors based on adenovirus type 5 and contain at least a functional part of the fiber protein from adenovirus type 16.

25 In another embodiment this invention provides adenoviral vectors or chimaeric adenoviral vectors that escape the liver following systemic administration. Preferably these adenoviral vectors are derived from subgroup A, B, D, or F in particular serotypes 12, 16, 28 and 40 or contain at

30 least the cell-binding moiety of the fiber protein derived from said adenoviruses.

It is to be understood that in all embodiments the adenoviral vectors may be derived from the serotype having the desired properties or that the adenoviral vector is

35 based on an adenovirus from one serotype and contains the sequences comprising the desired functions of another

serotype, these sequences replacing the native sequences in the said serotype.

- In another aspect this invention describes chimaeric adenoviruses and methods to generate these viruses that have an altered tropism different from that of adenovirus serotype 5. For example, viruses based on adenovirus serotype 5 but displaying any adenovirus fiber existing in nature. This chimaeric adenovirus serotype 5 is able to infect certain cell types more efficiently, or less efficiently *in vitro* and *in vivo* than the adenovirus serotype 5. Such cells include but are not limited to endothelial cells, smooth muscle cells, dendritic cells, neuronal cells, glial cells, synovial cells, lung epithelial cells, hemopoietic stem cells, monocytic/macrophage cells, tumor cells, skeletal muscle cells, mesothelial cells, synoviocytes, etc.
- 20 In another aspect the invention describes the construction and use of libraries consisting of distinct parts of adenovirus serotype 5 in which one or more genes or sequences have been replaced with DNA derived from alternative human or animal serotypes. This set of
- 25 constructs, in total encompassing the complete adenovirus genome, allows for the construction of unique chimaeric adenoviruses customized for a certain disease, group of patients or even a single individual.
- In all aspects of the invention the chimaeric adenoviruses may, or may not, contain deletions in the E1 region and insertions of heterologous genes linked either or not to a promoter. Furthermore, chimaeric adenoviruses may, or may not, contain deletions in the E3 region and insertions of heterologous genes linked to a promoter. Furthermore,
- 30 chimaeric adenoviruses may, or may not, contain deletions in the E2 and/or E4 region and insertions of heterologous genes

linked to a promoter. In the latter case E2 and/or E4 complementing cell lines are required to generate recombinant adenoviruses. In fact any gene in the genome of the viral vector can be taken out and supplied in trans.

5 Thus, in the extreme situation, chimaeric viruses do not contain any adenoviral genes in their genome and are by definition minimal adenoviral vectors. In this case all adenoviral functions are supplied in trans using stable cell lines and/or transient expression of these genes. A method

10 for producing minimal adenoviral vectors is described in WO97/00326 and is taken as reference herein. In another case Ad/AAV chimaeric molecules are packaged into the adenovirus capsids of the invention. A method for producing Ad/AAV chimaeric vectors is described in EP 97204085.1 and is taken

15 as reference herein. In principle any nucleic acid may be provided with the adenovirus capsids of the invention.

In one embodiment the invention provides a gene delivery vehicle having been provided with at least a tissue tropism

20 for smooth muscle cells and/or endothelial cells. In another embodiment the invention provides a gene delivery vehicle deprived of a tissue tropism for at least liver cells. Preferably, said gene delivery vehicle is provided with a tissue tropism for at least smooth muscle cells and/or

25 endothelial cells and deprived of a tissue tropism for at least liver cells. In a preferred embodiment of the invention said gene delivery vehicle is provided with a tissue tropism for at least smooth muscle cells and/or endothelial cells and/or deprived of a tissue tropism for at least liver cells

30 using a fiber protein derived from a subgroup B adenovirus, preferably of adenovirus 16. In a preferred aspect of the invention said gene delivery vehicle comprises a virus capsid. Preferably said virus capsid comprises a virus capsid derived in whole or in part from an adenovirus of subgroup B,

35 preferably from adenovirus 16, or it comprises proteins, or parts thereof, from an adenovirus of subgroup B, preferably of adenovirus 16. In preferred embodiment of the invention

said virus capsid comprises proteins, or fragments thereof, from at least two different viruses, preferably adenoviruses. In a preferred embodiment of this aspect of the invention at least one of said virus is an adenovirus of subgroup B, preferably adenovirus 16.

5 In a preferred embodiment of the invention said gene delivery vehicle comprises an adenovirus fiber protein or fragments thereof. Said fiber protein is preferably derived from an adenovirus of subgroup B, preferably of adenovirus 16. Said gene delivery vehicle may further comprise other fiber proteins, or fragments thereof, from other adenoviruses. Said gene delivery vehicle may or may not comprise other adenovirus proteins. Nucleic acid may be linked directly to fiber proteins, or fragments thereof, but may also be linked indirectly. Examples of indirect linkages include but are not limited to, packaging of nucleic acid into adenovirus capsids or packaging of nucleic acid into liposomes, wherein a fiber protein, or a fragment thereof, is incorporated into an adenovirus capsid or linked to a liposome. Direct linkage of nucleic acid to a fiber protein, or a fragment thereof, may be performed when said fiber protein, or a fragment thereof, is not part of a complex or when said fiber protein, or a fragment thereof, is part of complex such as an adenovirus capsid.

25 In one embodiment of the invention is provided a gene delivery vehicle comprising an adenovirus fiber protein wherein said fiber protein comprises a tissue determining fragment of an adenovirus of subgroup B adenovirus preferably of adenovirus 16. Adenovirus fiber protein comprises three functional domains. One domain, the base, is responsible for anchoring the fiber to a penton base of the adenovirus capsid. Another domain, the knob, is responsible for receptor recognition whereas the shaft domain functions as a spacer separating the base from the knob. The different domains may also have other function. For instance, the shaft is presumably also involved in target cell specificity. Each of the domains mentioned above may be used to define a fragment

of a fiber. However, fragments may also be identified in another way. For instance the knob domain comprises of a receptor binding fragment and a shaft binding fragment. The base domain comprises of a penton base binding fragment and a shaft binding fragment. Moreover, the shaft comprises of repeated stretches of amino acids. Each of these repeated stretches may be a fragment.

A tissue tropism determining fragment of a fiber protein may be a single fragment of a fiber protein or a combination of fragments of at least one fiber protein, wherein said tissue tropism determining fragment, either alone or in combination with a virus capsid, determines the efficiency with which a gene delivery vehicle can transduce a given cell or cell type, preferably but not necessarily in a positive way. With a tissue tropism for liver cells is meant a tissue tropism for cells residing in the liver, preferably liver parenchyma cells.

A tissue tropism for a certain tissue may be provided by increasing the efficiency with which cells of said tissue are transduced, alternatively, a tissue tropism for a certain tissue may be provided by decreasing the efficiency with which other cells than the cells of said tissue are transduced.

25

Fiber proteins possess tissue tropism determining properties. The most well described fragment of the fiber protein involved in tissue tropism is the knob domain. However, the shaft domain of the fiber protein also possesses tissue tropism determining properties. However, not all of the tissue tropism determining properties of an adenovirus capsid are incorporated into a fiber protein.

In a preferred embodiment of the invention, a fiber protein derived from a subgroup B adenovirus, preferably adenovirus 16, is combined with the non-fiber capsid proteins from an adenovirus of subgroup C, preferably of adenovirus 5.

In one aspect of the invention is provided a gene delivery vehicle comprising a nucleic acid derived from an adenovirus. In a preferred embodiment of the invention said adenovirus nucleic acid comprises at least one nucleic acid sequence  
5 encoding a fiber protein comprising at least a tissue tropism determining fragment of a subgroup B adenovirus fiber protein, preferably of adenovirus 16. In a preferred aspect said adenovirus comprises nucleic acid from at least two different adenoviruses. In a preferred aspect said adenovirus  
10 comprises nucleic acid from at least two different adenoviruses wherein at least one nucleic acid sequence encoding a fiber protein comprising at least a tissue tropism determining fragment of a subgroup B adenovirus fiber protein, preferably of adenovirus 16.

15 In a preferred embodiment of the invention said adenovirus nucleic acid is modified such that the capacity of said adenovirus nucleic acid to replicate in a target cell has been reduced or disabled. This may be achieved through inactivating or deleting genes encoding early region 1  
20 proteins.

In another preferred embodiment said adenovirus nucleic acid is modified such that the capacity of a host immune system to mount an immune response against adenovirus proteins encoded by said adenovirus nucleic acid has been reduced or disabled.  
25 This may be achieved through deletion of genes encoding proteins of early region 2 and/or early region 4. Alternatively, genes encoding early region 3 proteins, may be deleted, or on the contrary, considering the anti-immune system function of some of the proteins encoded by the genes  
30 in early region 3, the expression of early region 3 proteins may be enhanced for some purposes. Also, the adenovirus nucleic acid may be altered by a combination of two or more of the specific alterations of the adenovirus nucleic acid mentioned above. It is clear that when essential genes are  
35 deleted from the adenovirus nucleic acid, the genes must be complemented in the cell that is going to produce the adenovirus nucleic acid, the adenovirus vector, the vehicle

or the chimaeric capsid. The adenovirus nucleic acid may also be modified such that the capacity of a host immune system to mount an immune response against adenovirus proteins encoded by said adenovirus nucleic acid has been reduced or disabled, in other ways then mentioned above, for instance through exchanging capsid proteins, or fragments thereof, by capsid proteins, or fragments thereof, from other serotypes for which humans do not have, or have low levels of, neutralizing antibodies. Another example of this is the exchange of genes encoding capsid proteins with the genes encoding for capsid proteins from other serotypes. Also capsid proteins, or fragments thereof, may be exchanged for other capsid proteins, or fragments thereof, for which individuals are not capable of, or have a low capacity of, raising an immune response against.

An adenovirus nucleic acid may be altered further or instead of one or more of the alterations mentioned above, by inactivating or deleting genes encoding adenovirus late proteins such as but not-limited to, hexon, penton, fiber and/or protein IX.

In a preferred embodiment of the invention all genes encoding adenovirus proteins are deleted from said adenovirus nucleic acid, turning said nucleic acid into a minimal adenovirus vector.

In another preferred embodiment of the invention said adenovirus nucleic acid is an Ad/AAV chimaeric vector, wherein at least the integration means of an adeno-associated virus (AAV) are incorporated into said adenovirus nucleic acid.

In a preferred embodiment of the invention, a vector or a nucleic acid, which may be one and the same or not, according to the invention further comprises at least one non-adenovirus gene. Preferably, at least one of said non-adenovirus gene is selected from the group of genes encoding: an apolipoprotein, a ceNOS, a herpes simplex virus thymidine kinase, an interleukin-3, an interleukin-1 $\alpha$ , an

(anti)angiogenesis protein such as angiostatin, an anti-proliferation protein, a vascular endothelial growth factor (VGAF), a basic fibroblast growth factor (bFGF), a hypoxia inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ), a PAI-1 or a smooth muscle  
5 cell anti-migration protein.

In another aspect, the invention provides a cell for the production of a gene delivery vehicle provided with at least a tissue tropism for smooth muscle cells and/or endothelial cells. In another aspect, the invention provides a cell for  
10 the production of a gene delivery vehicle deprived of at least a tissue tropism for liver cells. In another aspect, the invention provides a cell for the production of a gene delivery vehicle provided with at least a tissue tropism for smooth muscle cells and/or endothelial cells and deprived of  
15 at least a tissue tropism for liver cells. In a preferred embodiment of the invention said cell is an adenovirus packaging cell, wherein an adenovirus nucleic acid is packaged into an adenovirus capsid. In one aspect of an adenovirus packaging cell of the invention all proteins  
20 required for the replication and packaging of an adenovirus nucleic acid, except for the proteins encoded by early region 1, are provided by genes incorporated in said adenovirus nucleic acid. The early region 1 encoded proteins in this aspect of the invention may be encoded by genes incorporated  
25 into the cells genomic DNA. In a preferred embodiment of the invention said cell is PER.C6 (ECACC deposit number 96022940). In general, when gene products required for the replication and packaging of adenovirus nucleic acid into adenovirus capsid are not provided by a adenovirus nucleic  
30 acid, they are provided by the packaging cell, either by transient transfection, or through stable transformation of said packaging cell. However, a gene product provided by the packaging cell may also be provided by a gene present on said adenovirus nucleic acid. For instance fiber protein may be  
35 provided by the packaging cell, for instance through transient transfection, and may be encoded by the adenovirus nucleic acid. This feature can among others be used to

generate adenovirus capsids comprising of fiber proteins from two different viruses.

The gene delivery vehicles of the invention are useful for the treatment cardiovascular disease or disease treatable by nucleic acid delivery to endothelial cells or smooth muscle cells. A non-limiting example of the latter is for instance cancer, where the nucleic acid transferred comprises a gene encoding an anti-angiogenesis protein.

The gene delivery vehicles of the invention may be used as a pharmaceutical for the treatment of said diseases.

Alternatively, gene delivery vehicles of the invention may be used for the preparation of a medicament for the treatment of said diseases.

In one aspect the invention provides an adenovirus capsid with or provided with a tissue tropism for smooth muscle cells and/or endothelial cells wherein said capsid preferably comprises proteins from at least two different adenoviruses and wherein at least a tissue tropism determining fragment of a fiber protein is derived from a subgroup B adenovirus, preferably of adenovirus 16. In another aspect the invention provides an adenovirus capsid deprived of a tissue tropism for liver cells wherein said capsid preferably comprises proteins from at least two different adenoviruses and wherein at least a tissue tropism determining fragment of a fiber protein is derived from a subgroup B adenovirus, preferably of adenovirus 16.

In one embodiment the invention comprises the use of an adenovirus capsid, for the delivery of nucleic acid to smooth muscle cells and/or endothelial cells. In another embodiment the invention comprises the use of an adenovirus capsid, for preventing the delivery of nucleic acid to liver cells.

The adenovirus capsids of the invention may be used for the treatment cardiovascular disease or disease treatable by nucleic acid delivery to endothelial cells or smooth muscle cells. Example of the latter is for instance cancer where the nucleic acid transferred comprises a gene encoding an anti-angiogenesis protein.

The adenovirus capsids of the invention may be used as a pharmaceutical for the treatment of said diseases.

Alternatively, adenovirus capsids of the invention may be used for the preparation of a medicament for the treatment of  
5 said diseases.

In another aspect of the invention is provided construct pBr/Ad.BamRAFib, comprising adenovirus 5 sequences 21562-31094 and 32794-35938.

In another aspect of the invention is provided construct  
10 pBr/AdBamRfib16, comprising adenovirus 5 sequences 21562-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein.

In yet another aspect of the invention is provided construct pBr/AdBamR.pac/fib16, comprising adenovirus 5 sequences  
15 21562-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein, and further comprising a unique PacI-site in the proximity of the adenovirus 5 right terminal repeat, in the non-adenovirus sequence backbone of said construct.

20 In another aspect of the invention is provided construct pWE/Ad.AflIIrITRfib16 comprising Ad5 sequence 3534-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein.

In another aspect of the invention is provided construct  
25 pWE/Ad.AflIIrITRDE2Afib16 comprising Ad5 sequences 3534-22443 and 24033-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein.

In the numbering of the sequences mentioned above, the number is depicted until and not until plus.

30

In a preferred embodiment of the invention said constructs are used for the generation of a gene delivery vehicle or an adenovirus capsid with a tissue tropism for smooth muscle cells and/or endothelial cells.

35 In another aspect the invention provides a library of adenovirus vectors, or gene delivery vehicles which may be one and the same or not, comprising a large selection of non-

adenovirus nucleic acids. In another aspect of the invention, adenovirus genes encoding capsid proteins are used to generate a library of adenovirus capsids comprising of proteins derived from at least two different adenoviruses, 5 said adenoviruses preferably being derived from two different serotypes, wherein preferably one serotype is an adenovirus of subgroup B. In a particularly preferred embodiment of the invention a library of adenovirus capsids is generated comprising proteins from at least two different adenoviruses 10 and wherein at least a tissue tropism determining fragment of fiber protein is derived from an adenovirus of subgroup B, preferably of adenovirus 16.

A fiber protein of adenovirus 16 preferably comprises of the 15 sequence given in figure 9. However within the scope of the present invention analogous sequences may be obtained through using codon degeneracy. Alternatively, amino-acid substitutions or insertions or deletions may be performed as long as the tissue tropism determining property is not 20 significantly altered. Such amino-acid substitutions may be within the same polarity group or without.

In the following the invention is illustrated by a number of non-limiting examples.

## EXAMPLES

### Example 1: Generation of adenovirus serotype 5 based viruses with chimaeric fiber proteins

#### 5    Generation of adenovirus template clones lacking DNA encoding for fiber

The fiber coding sequence of adenovirus serotype 5 is located between nucleotides 31042 and 32787. To remove the adenovirus serotype 5 DNA encoding fiber we started with construct

10    pBr/Ad.Bam-rITR (Figure 1; ECACC deposit P97082122). From this construct first a NdeI site was removed. For this purpose, pBr322 plasmid DNA was digested with NdeI after which protruding ends were filled using Klenow enzyme. This pBr322 plasmid was then re-ligated, digested with NdeI and

15    transformed into *E. coli* DH5 $\alpha$ . The obtained pBr/ $\Delta$ NdeI plasmid was digested with ScaI and SalI and the resulting 3198 bp vector fragment was ligated to the 15349 bp ScaI-SalI fragment derived from pBr/Ad.BamrITR, resulting in plasmid pBr/Ad.Bam-rITR $\Delta$ NdeI which hence contained a unique NdeI

20    site. Next a PCR was performed with oligonucleotides "NY-up" and "NY-down" (Figure 2). During amplification, both a NdeI and a NsiI restriction site were introduced to facilitate cloning of the amplified fiber DNAs. Amplification consisted of 25 cycles of each 45 sec. at 94°C, 1 min. at 60°C, and 45

25    sec. at 72°C. The PCR reaction contained 25 pmol of oligonucleotides NY-up or NY-down, 2mM dNTP, PCR buffer with 1.5 mM MgCl<sub>2</sub>, and 1 unit of Elongase heat stable polymerase (Gibco, The Netherlands). One-tenth of the PCR product was run on an agarose gel which demonstrated that the expected

30    DNA fragment of  $\pm$  2200 bp was amplified. This PCR fragment was subsequently purified using Geneclean kit system. (Bio101 Inc.) Then, both the construct pBr/Ad.Bam-rITR $\Delta$ NdeI as well as the PCR product were digested with restriction enzymes NdeI and SbfI. The PCR fragment was subsequently cloned using

T4 ligase enzyme into the NdeI and SbfI sites thus generating pBr/Ad.BamRAFib (Figure 3).

Amplification of fiber sequences from adenovirus serotypes

5 To enable amplification of the DNAs encoding fiber protein derived from alternative serotypes degenerate oligonucleotides were synthesized. For this purpose, first known DNA sequences encoding for fiber protein of alternative serotypes were aligned to identify conserved regions in both  
10 the tail region as well as the knob region of the fiber protein. From the alignment, which contained the nucleotide sequence of 19 different serotypes representing all 6 subgroups, (degenerate) oligonucleotides were synthesized (see Table I). Also shown in table 3 is the combination of  
15 oligonucleotides used to amplify the DNA encoding fiber protein of a specific serotype. The amplification reaction (50 µl) contained 2 mM dNTPs, 25 pmol of each oligonucleotide, standard 1x PCR buffer, 1,5 mM MgCl<sub>2</sub>, and 1 Unit Pwo heat stable polymerase (Boehringer Mannheim) per  
20 reaction. The cycler program contained 20 cycles, each consisting of 30 sec. 94°C, 60 sec. 60-64°C, and 120 sec. 72°C. One-tenth of the PCR product was run on an agarose gel to demonstrate that a DNA fragment was amplified. Of each different template, two independent PCR reactions were  
25 performed.

Generation of chimaeric adenoviral DNA constructs

All amplified fiber DNAs as well as the vector (pBr/Ad.BamRAFib) were digested with NdeI and NsiI. The  
30 digested DNAs were subsequently run on a agarose gel after which the fragments were isolated from the gel and purified using the Geneclean kit (Bio101 Inc). The PCR fragments were then cloned into the NdeI and NsiI sites of pBr/AdBamRAFib, thus generating pBr/AdBamRFibXX (where XX stands for the

serotype number of which the fiber DNA was isolated). The inserts generated by PCR were sequenced to confirm correct amplification. The obtained sequences of the different fiber genes are shown in Figure 4.

5

Generation of recombinant adenovirus chimaeric for fiber protein

To enable efficient generation of chimaeric viruses an AvrII fragment from the pBr/AdBamRFib16, pBr/AdBamRFib28, 10 pBr/AdBamRFib40-L constructs was subcloned into the vector pBr/Ad.Bam-rITR.pac#8 (ECACC deposit #P97082121) replacing the corresponding sequences in this vector. pBr/Ad.Bam-rITR.pac#8 has the same adenoviral insert as pBr/Ad.Bam-rITR but has a PacI site near the rITR that enables the ITR to be 15 separated from the vector sequences. The construct pWE/Ad.AflIII-Eco was generated as follows. PWE.pac was digested with ClaI and the 5 prime protruding ends were filled in with Klenow enzyme. The DNA was then digested with PacI and isolate from agarose gel. PWE/AflIIrITR was 20 digested with EcoRI and after treatment with Klenow enzyme digested with PacI. The large 24 kb. fragment containing the adenoviral sequences was isolated from agarose gel and ligated to the ClaI digested and blunted pWE.Pac vector. Use was made of the ligation express kit from Clontech. After 25 transformation of XL10-gold cells from Stratagene, clones were identified that contained the expected construct. PWE/Ad.AlfII-Eco contains Ad5 sequences from basepairs 3534-27336. Three constructs, pClipsal-Luc (Figure 5) digested with SalI, pWE/Ad.AflIII-Eco digested with PacI and EcoRI and 30 pBr/AdBamR.pac/fibXX digested with BamHI and PacI were transfected into adenovirus producer cells (PER.C6, Fallaux et al, 1998). Figure 6 schematically depicts the method and fragments used to generate the chimaeric viruses. Only pBr/Ad.BamRfib12 was used without subcloning in the PacI 35 containing vector and therefor was not liberated from vector sequences using PacI but was digested with ClaI which leaves

approximately 160 bp of vector sequences attached to the right ITR. Furthermore, the pBr/Ad.BamRfib12 and pBr/Ad.BamRfib28 contain an internal BamHI site in the fiber sequences and were therefor digested with SalI which cuts in the vector sequences flanking the BamHI site. For transfection, 2 µg of pCLIPsal-Luc, and 4 µg of both pWE/Ad.AflII-Eco and pBr/AdBamR.pac/fibXX were diluted in serum free DMEM to 100 µl total volume. To this DNA suspension 100 µl 2.5x diluted lipofectamine (Gibco) in serum-free medium was added. After 30 minutes at room temperature the DNA-lipofectamine complex solution was added to 2.5 ml of serum-free DMEM which was subsequently added to a T25 cm<sup>2</sup> tissue culture flask. This flask contained PER.C6 cells that were seeded 24-hours prior to transfection at a density of 1x10<sup>6</sup> cells/flask. Two hours later, the DNA-lipofectamine complex containing medium was diluted once by the addition of 2.5 ml DMEM supplemented with 20% fetal calf serum. Again 24 hours later the medium was replaced by fresh DMEM supplemented with 10% fetal calf serum. Cells were cultured for 6-8 days, subsequently harvested, and freeze/thawed 3 times. Cellular debris was removed by centrifugation for 5 minutes at 3000 rpm room temperature. Of the supernatant (12.5 ml) 3-5 ml was used to infect again PER.C6 cells (T80 cm<sup>2</sup> tissue culture flasks). This re-infection results in full cytopathogenic effect (CPE) after 5-6 days after which the adenovirus is harvested as described above.

#### Production of fiber chimeric adenovirus

10 ml of the above described crude lysate was used to inoculate a 1 liter fermentor which contained 1 - 1.5 x 10<sup>6</sup> PER.C6 cells/ml growing in suspension. Three days after inoculation, the cells were harvested and pelleted by centrifuging for 10 min at 1750 rpm at room temperature. The chimeric adenovirus present in the pelleted cells was subsequently extracted and purified using the following

downstream processing protocol. The pellet was dissolved in 50 ml 10 mM  $\text{NaPO}_4^-$  and frozen at  $-20^\circ\text{C}$ . After thawing at  $37^\circ\text{C}$ , 5.6 ml deoxycholate (5% w/v) was added after which the solution was homogenated. The solution was subsequently

5 incubated for 15 minutes at  $37^\circ\text{C}$  to completely crack the cells. After homogenizing the solution, 1875  $\mu\text{l}$  (1M)  $\text{MgCl}_2$  was added and 5 ml 100% glycerol. After the addition of 375  $\mu\text{l}$  DNase (10 mg/ ml) the solution was incubated for 30 minutes at  $37^\circ\text{C}$ . Cell debris was removed by centrifugation at

10 1880xg for 30 minutes at room temperature without the brake on. The supernatant was subsequently purified from proteins by loading on 10 ml of freon. Upon centrifugation for 15 minutes at 2000 rpm without brake at room temperature three bands are visible of which the upper band represents the

15 adenovirus. This band was isolated by pipetting after which it was loaded on a Tris/HCl (1M) buffered caesiumchloride blockgradient (range: 1.2 to 1.4 gr./ml). Upon centrifugation at 21000 rpm for 2.5 hours at  $10^\circ\text{C}$  the virus was purified from remaining protein and cell debris since the virus, in

20 contrast to the other components, does not migrate into the 1.4 gr./ ml caesiumchloride solution. The virus band is isolated after which a second purification using a Tris/ HCl (1M) buffered continuous gradient of 1.33 gr. /ml of caesiumchloride is performed. After virus loading on top of

25 this gradient the virus is centrifuged for 17 hours at 55000 rpm at  $10^\circ\text{C}$ . Subsequently the virus band is isolated and after the addition of 30  $\mu\text{l}$  of sucrose (50 w/v) excess caesiumchloride is removed by three rounds of dialysis, each round comprising of 1 hour. For dialysis the virus is

30 transferred to dialysis slides (Slide-a-lizer, cut off 10000 kDa, Pierce, USA). The buffers used for dialysis are PBS which are supplemented with an increasing concentration of sucrose (round 1 to 3: 30 ml, 60 ml, and 150 ml sucrose (50% w/v)/ 1.5 liter PBS, all supplemented with 7.5 ml 2% (w/v)

35  $\text{CaMgCl}_2$ ). After dialysis, the virus is removed from the slide-a-lizer after which it is aliquoted in portions of 25

and 100  $\mu$ l upon which the virus is stored at  $-85^{\circ}\text{C}$ . To determine the number of virusparticles per ml, 50  $\mu$ l of the virus batch is run on an high pressure liquid chromatograph (HPLC) as described by Shamram et al (1997). The virus titers were found to be in the same range as the Ad5.Luc virus batch (Ad5.Luc:  $2.2 \times 10^{11}$  vp/ ml; Ad5.LucFib12:  $1.3 \times 10^{11}$  vp/ ml; Ad5.LucFib16:  $3.1 \times 10^{12}$  vp/ ml; Ad5.LucFib28:  $5.4 \times 10^{10}$  vp/ ml; Ad5.LucFib40-L:  $1.6 \times 10^{12}$  vp/ ml).

10 **Example 2: biodistribution of chimeric viruses after intravenous tail vein injection of rats.**

To investigate the biodistribution of the chimeric adenoviruses carrying fiber 12, 16, 28, or 40-2,  $1 \times 10^{10}$  particles of each of the generated virusbatches was diluted in 1 ml PBS after which the virus was injected in the tail vein of adult male Wag/Rij rats (3 rats/ virus). As a control, Ad5 carrying the luciferase transgene was used. Forty-eight hours after the administration of the virus, the rats were sacrificed after which the liver, spleen, lung, kidney, heart, and brain were dissected. These organs were subsequently mixed with 1 ml of lysis buffer (1% Triton X-100/ PBS) and minced for 30 seconds to obtain a protein lysate. The protein lysate was subsequently tested for the presence of transgene expression (luciferase activity) and the protein concentration was determined to express the luciferase activity per  $\mu$ g of protein. The results, Shown in Table II, demonstrate that in contrast to the Adenovirus serotype 5 control, none of the fiber chimeras are targeted specifically to the liver or to the spleen. This experiment shows that it is possible to circumvent the uptake of adenoviruses by the liver by making use of fibers of other serotypes. It also shows that the uptake by the liver is not correlated with the length of the fiber shaft, or determined solely by the ability of fiber knob to bind to CAR. The fibers used have different shaft lengths and, except for

fiber 16, are derived from subgroups known to have a fiber that can bind CAR (Roelvink et al 1998).

**Example 3: Chimeric viruses display differences in  
5 endothelial and smooth muscle cell transduction**

**A) Infection of Human endothelial cells**

Human endothelial cells (HUVEC) were isolated, cultured and  
10 characterized as described previously (Jaffe et al 1973;  
Wijnberg et al 1997). Briefly, cells were cultured on  
gelatin-coated dishes in M199 supplemented with 20 mM HEPES,  
pH 7.3 (Flow Labs., Irvine, Scotland), 10% (v/v) human serum  
(local blood bank), 10% (v/v) heat-inactivated newborn calf  
15 serum (NBCS) (GIBCO BRL, Gaithersburg, MD), 150 µg/ ml crude  
endothelial cell growth factor, 5 U/ ml heparin (Leo  
Pharmaceutics Products, Weesp, The Netherlands), penicillin  
(100 IU/ ml)/streptomycin (100 µg/ ml) (Boehringer Mannheim,  
Mannheim, FRG) at 37°C under 5% (v/v) CO<sub>2</sub>/ 95% (v/v) air  
20 atmosphere. Cells used for experiments were between passage  
1-3. In a first set of experiments 40000 HUVEC cells (a pool  
from 4 different individuals) were seeded in each well of 24-  
wells plates in a total volume of 200 µl. Twenty-four hours  
after seeding, the cells were washed with PBS after which 200  
25 µl of DMEM supplemented with 2% FCS was added to the cells.  
This medium contained various amounts of virus (MOI = 50,  
250, 1000, 2500, 5000, and 10000). The viruses used were  
besides the control Ad5, the fiber chimeras 12, 16, 28, and  
40-L (each infection in triplicate). Two hours after addition  
30 of the virus the medium was replaced by normal medium. Again  
forty-eight hours later cells were washed and lysed by the  
addition of 100 µl lysisbuffer. In figure 7a, results are  
shown on the transgene expression per microgram total protein  
after infection of HUVEC cells. These results show that fiber  
35 chimeras 12 and 28 are unable to infect HUVEC cells, that 40-  
L infects HUVECs with similar efficiency as the control Ad5

virus, and that fiber chimera 16 infects HUVECs significantly better. In a next set of experiments ( $n = 8$ ) the fiber 16 chimera was compared with the Ad5.Luc vector on HUVEC for luciferase activity after transduction with 2500 virus particles per cell of each virus. These experiments demonstrated that fiber 16 yields, on average, 8.1 fold increased luciferase activity ( $SD \pm 4.6$ ) as compared with Ad5. In a next experiment, an equal number of virus particles was added to wells of 24-well plates that contained different HUVEC cell concentrations. This experiment was performed since it is known that HUVECs are less efficiently infected with adenovirus serotype 5 when these cells reach confluency. For this purpose, HUVECs were seeded at 22500, 45000, 90000, and 135000 cells per well of 24-well plates (in triplicate). Twenty-four hours later these cells were infected as described above with medium containing  $4.5 \times 10^8$  virusparticles. The viruses used were, besides the control adenovirus serotype 5, the chimera fiber 16. The result of the transgene expression (RLU) per microgram protein determined 48 hours after infection (see figure 7b) shows that the fiber 16 chimeric adenovirus is also better suited to infect HUVEC cells even when these cells are 100% confluent which better mimics an *in vivo* situation. Since the Luciferase markergene does not provide information concerning the number of cells infected another experiment was performed with adenovirus serotype 5 and the fiber 16 chimera, both carrying a green fluorescent protein (GFP) as a marker gene. This protein expression can be detected using a flow cytometer which renders information about the percentage of cells transduced as well as fluorescence per cell. In this experiment cells were seeded at a concentration of 40000 cells per well and were exposed to virus for 2 hours. The virus used was Ad5.GFP ( $8.4 \times 10^{11}$  vp/ ml) and Ad5.Fib16.GFP ( $5.1 \times 10^{11}$  vp/ ml). Cells were exposed to a virus concentration of 500 virus particles per cell. Flow cytometric analysis, 48 hours after virus exposure

demonstrated that the fiber 16 virus gives higher transgene expression levels per cell since the median fluorescence, a parameter identifying the amount of GFP expression per cell, is higher with fiber 16 as compared to Ad5 (Figure 7c). These results are thus consistent and demonstrate that the fiber 16 chimeric virus is better suited to infect human primary endothelial cells as compared to Ad5.

#### B) Infection of human smooth muscle cells

Smooth muscle cells were isolated after isolation of EC (Weinberg et al 1997). The veins were incubated with medium (DMEM) supplemented with penicillin/ streptomycin) containing 0.075% (w/v) collagenase (Worthington Biochemical Corp., Freehold, NJ, USA). After 45 minutes the incubation medium containing detached cells was flushed from the veins. Cells were washed and cultured on gelatin coated dishes in culture medium supplemented with 10% fetal calf serum and 10% human serum at 37°C under 5% (v/v) CO<sub>2</sub>/ 95% (v/v) air atmosphere. Cells used for experiments were between passage 3-6. We first tested the panel of chimeric fiber viruses versus the control adenovirus serotype 5 for the infection of human smooth muscle cells. For this purpose, 40000 human umbilical vein smooth muscle cells (HUVsmc) were seeded in wells of 24-well plates in a total volume of 200 µl. Twenty-four hours after seeding, the cells were washed with PBS after which 200 µl of DMEM supplemented with 2% FCS was added to the cells. This medium contained various amounts of virus (MOI = 50, 250, 1250, 2500, and 5000). The viruses used were besides the control Ad5 the fiber chimeras 12, 16, 28, and 40-L (each infection in triplicate). Two hours after addition of the virus the medium was replaced by normal medium. Again forty-eight hours later cells were washed and lysed by the addition of 100 µl lysisbuffer. In figure 8a, results are shown of the transgene expression per microgram total protein after infection of HUVsmc cells. These results show that fiber

chimeras 12 and 28 are unable to infect HUVsmc cells, that 40-L infects HUVsmc with similar efficiency as the control Ad5 virus, and that fiber chimera 16 infects HUVsmc significantly better. In a next set of experiments, smooth muscle cells derived from saphenous vene, arteria Iliaca, left interior mammary artery (LIMA) and aorta were tested for infection with the fiber 16 chimera and Ad5 (both carrying luciferase as a marker gene). These experiments (n = 11) demonstrated that, on average, the fiber 16 chimera yielded 61.4 fold increased levels in luciferase activity (SD  $\pm$  54.8) as compared to Ad5. The high standard deviation (SD) is obtained due to the finding that the adenoviruses used vary in their efficiency of infection of SMC derived from different human vessels. In a next experiment, an equal number of virus particles was added to wells of 24-well plates that contained different HUVsmc cell concentrations confluency. For this purpose, HUVsmc were seeded at 10000, 20000, 40000, 60000, and 80000 cells per well of 24-well plates (in triplicate). Twenty-four hours later these cells were infected as described above with medium containing  $2 \times 10^8$  virusparticles. The viruses used were, besides the control adenovirus serotype 5, the chimera fiber 16. The result of the transgene expression (RLU) per microgram protein determined 48 hours after infection (see figure 8b) shows that the fiber 16 chimeric adenovirus is better suited to infect smooth muscle cells even when these cells are 100% confluent which better mimics an *in vivo* situation.

To identify the number of SMCs transduced with the fiber 16 chimera and Ad5, we performed transduction experiments with Ad5.GFP and Ad5Fib16.GFP (identical batches as used for EC infections). Human umbilical vein SMC were seeded at a concentration of 60000 cells per well in 24-well plates and exposed for 2 hours to 500 or 5000 virus particles per cell of Ad5.GFP or Ad5Fib16.GFP. Forty-eight hours after exposure cells were harvested and analyzed using a flow cytometer. The results obtained show that the fiber 16 mutant yields

approximately 10 fold higher transduction of SMC since the GFP expression measured after transduction with 5000 virus particles of Ad5.GFP is equal to GFP expression after transduction with 500 virus particles per cell of the fiber 16 chimera (Figure 8c).

#### C) Subgroup B fiber mutants other than fiber 16

The shaft and knob of fiber 16 are derived from adenovirus serotype 16 which, as described earlier, belongs to subgroup B. Based on hemagglutination assays, DNA restriction patterns, and neutralization assays the subgroup B viruses have been further subdivided into subgroup B1 and B2 (Wadell et al 1984). Subgroup B1 members include serotypes 3, 7, 16, 21, and 51. Subgroup B2 members include 11, 14, 34, and 35. To test whether the increased infection of smooth muscle cells is a trade of all fibers derived from subgroup B or specific for one or more subgroup B fiber molecules, we compared fiber 16 and fiber 51 (both subgroup B1) with fiber 11 and fiber 35 (both subgroup B2). For this purpose HUVsmc were infected with increasing amounts of virus particles per cell (156, 312, 625, 1250, 2500, 5000). The fiber mutant all carry the Luciferase marker gene (Ad5Fib11.Luc:  $1.1 \times 10^{12}$  vp/ml; Ad5Fib35Luc:  $1.4 \times 10^{12}$  vp/ml ; Ad5Fib51Luc:  $1.0 \times 10^{12}$  vp/ml). Based on the Luciferase activity measured and shown in Figure 8d, efficient infection of SMC is not a general trade of all subgroup B fiber molecules. Clearly fiber 16 and fiber 11 are better suited for infection of SMC than fiber 35 and fiber 51. Nevertheless, all subgroup B fiber mutants tested infect SMC better as compared to Ad5.

#### D) Organ culture experiments

We next identified whether the observed difference in transduction of EC and SMC using the fiber 16 chimera or the Ad5 can also be demonstrated in organ culture experiments. Hereto, We focused on the following tissues: 1) Human

Saphenous vein: the vein used in approximately 80% of all clinical vein grafting procedures

2) Human pericard/ epicard: for delivery of recombinant adenoviruses to the pericardial fluid which after infection of the pericardial or epicardial cells produce the protein of interest from the transgene carried by the adenovirus.

3) Human coronary arteries: for percutaneous transluminal coronary angioplasty (PTCA) to prevent restenosis. Of the coronary arteries we focused on the Left artery descending (LAD) and right coronary artery (RCA).

Parts of a human saphenous vein left over after vein graft surgery were sliced into pieces of approximately 0.5 cm. These pieces (n=3) were subsequently cultured for 2 hours in 200 ml of  $5 \times 10^{10}$  virus particles per ml. After two hours virus exposure the pieces were washed with PBS and cultured for another 48 hours at 37°C in a 10% CO<sub>2</sub> incubator. The pieces were then washed, fixated and stained for LacZ transgene expression. The viruses were Ad5.LacZ ( $2.2 \times 10^{12}$  vp/ ml), the fiber 16 chimera Ad5Fib16.LacZ ( $5.2 \times 10^{11}$  vp/ ml), and A fiber 51 chimera: Ad5Fib51.LacZ ( $2.1 \times 10^{12}$  vp/ ml). The pieces of saphenous vein were macroscopically photographed using a digital camera. Based on LacZ transgene expression obtained after 2 hours of virus exposure on saphenous vein slices, both the fiber 16 and the fiber 51 chimeric viruses give higher infection since much more blue staining is observed using these viruses as compared to Ad5.LacZ (Figure 8e). Identical experiments as described on saphenous vein were performed with human pericard and the human coronary arteries: RCA and LAD. Results of these experiments (Figures 8f-8g-8h respectively) together with the experiments performed on primary cells confirmed the superiority of the fiber 16 and 51 mutants as compared to Ad5 in infecting human cardiovascular tissues.

E) CAR and integrin expression on human EC and SMC

From the above described results it is clear that the chimeric adenovirus with the shaft and knob from fiber 16 is well suited to infect endothelial cells and smooth muscle cells. Thus, by changing the fiber protein on Ad5 viruses we are able to increase infection of cells that are poorly infected by Ad5. The difference between Ad5 and Ad5Fib16, although significant on both cell types, is less striking on endothelial cells as compared to smooth muscle cells. This may reflect differences in receptor expression. For example, HUVsmc significantly more  $\alpha_v\beta_5$  integrins than HUVEC (see below). Alternatively, this difference may be due to differences in expression of the receptor of fiber 16. Ad5.LucFib16 infects umbilical vein smooth muscle cells approximately 8 fold better than umbilical vein endothelial cells whereas in case of Ad5.Luc viruses endothelial cells are better infected than smooth muscle cells. To test whether Ad5 infection correlated with receptor expression of these cells the presence of CAR and  $\alpha_v$ -integrins was assayed on a flow cytometer. For this purpose  $1 \times 10^5$  HUVEC cells or HUVsmc were washed once with PBS/ 0.5% BSA after which the cells were pelleted by centrifugation for 5 minutes at 1750 rpm at room temperature. Subsequently, 10  $\mu$ l of a 100 times diluted  $\alpha_v\beta_3$  antibody (Mab 1961, Brunswick chemie, Amsterdam, The Netherlands), a 100 times diluted antibody  $\alpha_v\beta_5$  (antibody Mab 1976, Brunswick chemie, Amsterdam, The Netherlands), or 2000 times diluted CAR antibody was a kind gift of Dr. Bergelson, Harvard Medical School, Boston, USA (Hsu et al) was added to the cell pellet after which the cells were incubated for 30 minutes at 4°C in a dark environment. After this incubation, cells were washed twice with PBS/0.5% BSA and again pelleted by centrifugation for 5 minutes at 1750 rpm room temperature. To label the cells, 10 ml of rat anti mouse IgG1 labeled with phycoerythrine (PE) was added to the cell pellet upon which the cells were again incubated for 30 minutes at 4°C in a dark environment. Finally the cells were washed twice with PBS/0.5% BSA and analyzed on a flow

cytometer. The results of these experiments are shown in table III. From the results it can be concluded that HUVsmc do not express detectable levels of CAR confirming that these cells are difficult to transduce with an adenovirus which enters the cells via the CAR receptor.

#### F) Infection of human A549 cells

As a control for the experiments performed on endothelial cells and smooth muscle cells, A549 cells were infected to establish whether an equal amount of virus particles of the different chimeric adenoviruses show significant differences in transgene expression on cell lines that are easily infected by adenovirus. This is to investigate whether the observed differences in infection efficiency on endothelial and smooth muscle cells are cell type specific. For this purpose,  $10^5$  A549 cells were seeded in 24-well plates in a volume of 200  $\mu$ l. Two hours after seeding the medium was replaced by medium containing different amounts of particles of either fiber chimera 5, 12, 16, or 40-L (MOI = 0, 5, 10, 25, 100, 500). Twenty-four hours after the addition of virus, the cells were washed once with PBS after which the cells were lysed by the addition of 100  $\mu$ l lysisbuffer to each well (1% Triton X-100 in PBS) after which transgene expression (Luciferase activity) and the protein concentration was determined. Subsequently, the luciferase activity per  $\mu$ g protein was calculated. The data, shown in table IV, demonstrate that Ad5.Luc viruses infect A549 cells most efficient while the infection efficiency of Ad5LucFib16 or Ad5LucFib40-L is a few times lower. This means that the efficient infection of endothelial cells and especially smooth muscle cells is due to differences in binding of the virus to these cells and not to the amount of virus or the quality of the viruses used.

Table I

	Serotype	Tail oligonucleotide	Knob oligonucleotide
5	4	A	1
	8	B	2
	9	B	2
	12	E	3
	16	C	4
10	19p	B	2
	28	B	2
	32	B	2
	36	B	2
	37	B	2
	40-1	D	5
	40-2	D	6
15	41-8	D	5
	41-1	D	7
	49	B	2
	50	B	2
20	51	C	8
25	A:	5'- CCC GTG TAT CCA TAT GAT GCA GAC AAC GAC CGA CC- 3'	
	B:	5'- CCC GTC TAC CCA TAT GGC TAC GCG CGG- 3'	
	C:	5'- CCK GTS TAC CCA TAT GAA GAT GAA AGC- 3'	
	D:	5'- CCC GTC TAC CCA TAT GAC ACC TYC TCA ACT C- 3'	
	E:	5'- CCC GTT TAC CCA TAT GAC CCA TTT GAC ACA TCA GAC- 3'	
	1:	5'- CCG ATG CAT TTA TTG TTG GGC TAT ATA GGA - 3'	
	2:	5'- CCG ATG CAT TYA TTC TTG GGC RAT ATA GGA - 3'	
	3:	5'- CCG ATG CAT TTA TTC TTG GGR AAT GTA WGA AAA GGA - 3'	
30	4:	5'- CCG ATG CAT TCA GTC ATC TTC TCT GAT ATA - 3'	
	5:	5'- CCG ATG CAT TTA TTG TTC AGT TAT GTA GCA - 3'	
	6:	5'- GCC ATG CAT TTA TTG TTC TGT TAC ATA AGA - 3'	
	7:	5'- CCG TTA ATT AAG CCC TTA TTG TTC TGT TAC ATA AGA A - 3'	
	8:	5'- CCG ATG CAT TCA GTC ATC YTC TWT AAT ATA - 3'	
35			

Table II

Organ	Control Ad5	Fib 12	Fib 16	Fib 28	Fib 40-L
Liver	740045	458	8844	419	2033
Spleen	105432	931	3442	592	16107
Lung	428	315	334	316	424
Kidney	254	142	190	209	224
Heart	474	473	276	304	302
Brain	291	318	294	323	257

Table III

Cell line	$\alpha_v\beta_3$	$\alpha_v\beta_5$	CAR
HUVEC 70%	98.3%	18.9%	18.1%
HUVEC 100%	97.2%	10.5%	7.2%
HUVsmc 70%	95.5%	76.6%	0.3%
HUVsmc 100%	92.2%	66.5%	0.3%
PER.C6	7.8%	16.8%	99.6%

Table IV

MOI (VP/Cell)	Control Ad5	Fiber 12	Fiber 16	Fiber 40-L
0	0	0	0	0
5	1025	46	661	443
10	1982	183	1704	843
25	4840	200	3274	2614
100	21875	1216	13432	11907
500	203834	3296	93163	71433

## REFERENCES

- Arnberg N., Mei Y. and Wadell G., 1997. Fiber genes of adenoviruses with tropism for the eye and the genital tract. *Virology* 227: 239-244.
- 5 Bergelson, J.M., Cunningham, J.A., Droguett, G., Kurt-Jones, E.A., Krithivas, A., Hong, J.S., Horwitz, M.S., Crowell, R.L. and Finberg, R.W. (1997) Isolation of a common receptor for coxsackie B virus and adenoviruses 2 and 5. *Science* 275: 1320-1323.
- 10 Bout A. (1997) Gene therapy, p. 167-182. In: D.J.A. Crommelin and R.D. Sindelar (ed.), *Pharmaceutical Biotechnology*, Harwood Academic Publishers.
- 15 Bout, A. (1996) Prospects for human gene therapy. *Eur. J. Drug Met. and Pharma.* 2, 175-179.
- Blaese, M., Blankenstein, T., Brenner, M., Cohen-Hagenauer, O., Gansbacher, B., Russel, S., Sorrentino, B. and Velu, T.
- 20 (1995) *Cancer Gene Ther.* 2: 291-297.
- Brody, S.L. and Crystal, R.G. (1994) Adenovirus mediated in vivo gene transfer. *Ann. N. Y. Acad. Sci.* 716:90-101.
- 25 Carter, A.J., Laird, J.R., Farb, A., Kufs, W., Wortham, D.C. and Virmani, R. (1994) Morphologic characteristics of lesion formation and time course of smooth muscle cell proliferation in a porcine proliferative restenosis model. *J. Am. Coll. Cardiol.* 24: 1398-1405.
- 30 Chroboczek J., Ruigrok R.W.H., and Cusack S., 1995. Adenovirus fiber, p. 163-200. In: W. Doerfler and P. Bohm (ed.), *The molecular repertoire of adenoviruses*, I. Springer-Verlag, Berlin.

- Defer C., Belin M., Caillet-Boudin M. and Boulanger P., 1990.  
Human adenovirus-host cell interactions; comparative study  
with members of subgroup B and C. *Journal of Virology* 64 (8):  
5 3661-3673.
- Fallaux, F.J., Bout, A., van der Velde, I et al. New helper  
cells and matched E1-deleted adenovirus vectors prevent  
generation of replication competent adenoviruses. *Human Gene*  
10 *Therapy*, 9 (1998), p1909-1917.
- Francki, R.I.B., Fauquet, C.M., Knudson, D.L. and Brown, F.  
(1991) Classification and nomenclature of viruses. Fifth  
report of the international Committee on taxonomy of viruses.  
15 *Arch. Virol. Suppl.* 2: 140-144.
- Gall J., Kass-Eisler A., Leinwand L. and Falck-Pedersen E.  
(1996) Adenovirus type 5 and 7 capsid chimera: fiber  
replacement alters receptor tropism without affecting primary  
20 immune neutralization epitopes. *Journal of Virology* 70 (4):  
2116-2123.
- Greber, U.F., Willets, M., Webster, P., and Helenius, A.  
(1993). Stepwise dismanteling of adenovirus 2 during entry  
25 into cells. *Cell* 75: 477-486.
- Hynes, R.O. (1992) Integrins: versatility, modulation and  
signalling in cell adhesion. *Cell* 69: 11-25.
- 30 Herz, J. and Gerard, R.D. (1993) Adenovirus-mediated transfer  
of low density lipoprotein receptor gene acutely accelerates  
cholesterol clearance in normal mice. *Proc. Natl. Acad. Sci.*  
*U.S.A.* 96: 2812-2816.
- 35 Hierholzer, J.C. (1992) Adenovirus in the immunocompromised  
host. *Clin. Microbiol Rev.* 5, 262-274.

Hierholzer, J.C., Wigand, R., Anderson, L.J., Adrian, T., and Gold, J.W.M. (1988) Adenoviruses from patients with AIDS: a plethora of serotypes and a description of five new serotypes  
5 of subgenus D (types 43-47). J. Infect. Dis. 158, 804-813.

Hong, S.S., Karayan, L., Tournier, J., Curiel, D.T. and Boulanger, P.A. (1997) Adenovirus type 5 fiber knob binds to MHC class I  $\alpha 2$  domain at the surface of human epithelial and  
10 B lymphoblastoid cells. EMBO J. 16: 2294-2306.

Hsu, K.H., Lonberg-Holm, K., Alstein, B. and Crowell, R.L. (1988)  
A monoclonal antibody specific for the cellular receptor for  
15 the group B coxsackieviruses. J. Virol 62(5): 1647-1652.

Huard, J., Lochmuller, H., Acsadi, G., Jani, A., Massie, B. and Karpati, G. (1995) The route of administration is a major determinant of the transduction efficiency of rat tissues by  
20 adenoviral recombinants. Gene Ther. 2: 107-115.

Ishibashi, M. and Yasue, H. (1984) The adenoviruses, H.S. Ginsberg, ed., Plenum Press, London, New York. Chapter 12, 497-561.

25 Jaffe, E.A., Nachman, R.L., Becker, C.G., Minick, C.R. (1973) Culture of endothelial cells derived from umbilical veins. Identification by morphologic and immunologic criteria. J. Clin. Invest. 52, 2745-2756.

30 Kass-Eisler, A., Falck-Pederson, E., Elfenbein, D.H., Alvira, M., Buttrick, P.M. and Leinwand, L.A. (1994) The impact of developmental stage, route of administration and the immune system on adenovirus-mediated gene transfer. Gene Ther. 1:  
35 395-402.

- Khoo, S.H., Bailey, A.S., De Jong, J.C., and Mandal, B.K. (1995). Adenovirus infections in human immunodeficiency virus-positive patients: Clinical features and molecular epidemiology. *J. Infect. Dis* 172, 629-637
- 5 Kidd, A.H., Chroboczek, J., Cusack, S., and Ruigrok, R.W. (1993) Adenovirus type 40 virions contain two distinct fibers. *Virology* 192, 73-84.
- 10 Krasnykh V.N., Mikheeva G.V., Douglas J.T. and Curiel D.T. (1996) Generation of recombinant adenovirus vectors with modified fibers for altering viral tropism. *J. Virol.* 70(10): 6839-6846.
- 15 Krasnykh V.N., Dmitriev I., Mikheeva G., Miller C.R., Belousova N. and Curiel D.T. (1998) Characterization of an adenovirus vector containing a heterologous peptide epitope in the HI loop of the fiber knob. *J. Virol.* 72(3): 1844-1852.
- 20 Law, L., Chillon, M., Bosch, A., Armentano, D., Welsh, M.J. and Davidson, B.L. (1998) Infection of primary CNS cells by different adenoviral serotypes: Searching for a more efficient vector. Abstract 1<sup>st</sup> Annual Meeting American Society of Gene Therapy, Seattle, Washington.
- 25 Leppard, K.N. (1997) E4 gene function in adenovirus, adenovirus vector and adeno-associated virus infections. *J. Gen. Virol.* 78: 2131-2138.
- 30 Lloyd Jones, D.M. and Bloch, K.D. (1996) The vascular biology of nitric oxide and its role in atherogenesis. *Annu. Rev. Med.* 47: 365-375.
- Morgan, C., Rozenkrantz, H.S., and Mednis, B. (1969)
- 35 Structure and development of viruses as observed in the electron microscope.X. Entry and uncoating of adenovirus. *J. Virol* 4, 777-796.

- Roelvink, P.W., Kovesdi, I. and Wickham, T.J. (1996)  
Comparative analysis of adenovirus fiber-cell interaction:  
Adenovirus type 2 (Ad2) and Ad9 utilize the same cellular  
5 fiber receptor but use different binding strategies for  
attachemnt. J. Virol. 70: 7614-7621.
- Roelvink, P.W., Lizonova, A., Lee, J.G.M., Li, Y., Bergelson,  
J.M., Finberg, R.W., Brough, D.E., Kovesdi, I. and Wickham,  
10 T.J. (1998) The coxsackie-adenovirus receptor protein can  
function as a cellular attachment protein for adenovirus  
serotypes from subgroups A, C, D, E, and F. J. Virol. 72:  
7909-7915.
- 15 Rogers, B.E., Douglas J.T., Ahlem, C., Buchsbaum, D.J.,  
Frincke, J. and Curiel, D.T. (1997) Use of a novel cross-  
linking method to modify adenovirus tropism. Gene Ther.4:  
1387-1392.
- 20 Schulick, A.H., Vassalli, G., Dunn, P.F., Dong, G., Rade,  
J.J., Zamarron, C. and Dichek, D.A. (1997). Established  
immunity precludes adenovirus-mediated gene transfer in rat  
carotid arteries.
- 25 Schnurr, D and Dondero, M.E. (1993) Two new candidate  
adenovirus serotypes. Intervirol. 36, 79-83.
- Schwartz, R.S., Edwards, W.D., Huber, K.C., Antoniudes, L.C.  
Bailey, K.R., Camrud, A.R., Jorgenson, M.A. and Holmes, D.R.  
30 Jr. (1993) Coronary restenosis: Prospects for solution and  
new perspectives from a porcine model. Mayo Clin. Proc. 68:  
54-62.
- Shi, Y., Pieniek, M., Fard, A., O'Brien, J., Mannion, J.D.  
35 and Zalewski, A. (1996) Adventitial remodelling after  
coronary arterial injury. Circulation 93: 340-348.

- Shabram, P.W., Giroux, D.D., Goudreau, A.M., Gregory, R.J., Horn, M.T., Huyghe, B.G., Liu, X., Nunnally, M.H., Sugarman, B.J. and Sutjipto, S. (1997) Analytical anion-exchange HPLC of recombinant type-5 adenoviral particles. Hum. Gene Ther. 8(4): 453-465.
- Signas, G., Akusjarvi, G., and Pettersson, U. (1985). Adenovirus 3 fiberpolypeptide gene: Complications for the structure of the fiber protein. J. Virol. 53, 672-678.
- Stevenson S.C., Rollence M., White B., Weaver L. and McClelland A., (1995) Human adenovirus serotypes 3 and 5 bind to two different cellular receptors via the fiber head domain. J. Virol 69(5): 2850-2857.
- Stevenson S.C., Rollence M., Marshall-Neff J. and McClelland A. (1997) Selective targeting of human cells by a chimaeric adenovirus vector containing a modified fiber protein. J. Virology 71(6): 4782-4790.
- Stouten, P.W.F., Sander, C., Ruigrok, R.W.H., and Cusack, S. (1992) New triple helical model for the shaft of the adenovirus fiber. J. Mol. Biol. 226, 1073-1084.
- Svensson, V. and Persson, R. (1984). Entry of adenovirus 2 into Hela cells. J. Virol. 51, 687-694.
- Van der Vliet, P.C. (1995) Adenovirus DNA replication In: W. Doerfler and P. Böhm (eds.), The molecular repertoire of adenoviruses II. Springer-Verlag, Berlin.
- Varga, M.J., Weibull, C., and Everitt, E. (1991). Infectious entry pathway of adenovirus type 2. J. Virol 65, 6061-6070.
- Varenne, O., Pislaru, S., Gillijns, H., Van Pelt, N., Gerard, R.D., Zoldhelyi, P., Van de Werf, F., Collen, D. and

- Janssens, S.P. (1998) Local adenovirus-mediated transfer of human endothelial nitric oxide synthetase reduces luminal narrowing after coronary angioplasty in pigs. *Circulation* 98: 919-926.
- 5 Wadell G (1984) Molecular Epidemiology of human adenoviruses *Curr. Top. Microbiol. Immunol.* 110, 191-220.
- 10 Wickham T.J., Carrion M.E. and Kovesdi I., 1995. Targeting of adenovirus penton base to new receptors through replacement of its RGD motif with other receptor-specific peptide motifs. *Gene Therapy* 2: 750-756.
- 15 Wickham T.J., Segal, D.M., Roelvink, P.W., Carrion M.E., Lizonova, A., Lee, G-M., and Kovesdi, I. (1996) Targeted adenovirus gene transfer to endothelial and smooth muscle cells by using bispecific antibodies. *J. Virol.* 70 (10), 6831-6838.
- 20 Wickham, T.J., Mathias, P., Cherish, D.A., and Nemerow, G.R. (1993) Integrins avb3 and avb5 promote adenovirus internalization but not virus attachment. *Cell* 73, 309-319.
- 25 Wijnberg, M.J., Quax, P.H.A., Nieuwenbroek, N.M.E., Verheijen, J.H. (1997). The migration of human smooth muscle cells in vitro is mediated by plasminogen activation and can be inhibited by  $\alpha_2$ - macro globulin receptor associated protein. *Thromb. and Haemostas.* 78, 880-886.
- 30 Wold, W.S., Tollefson, A.E. and Hermiston, T.W. (1995) E3 transcription unit of adenovirus. In: W. Doerfler and P. Böhm (eds.), *The molecular repertoire of adenoviruses I*. Springer-Verlag, Berlin.

Zabner, J., Armentano, D., Chillon, M., Wadsworth, S.C. and Welsh, M.J. (1998) Type 17 fiber enhances gene transfer Abstract 1<sup>st</sup> Annual Meeting American Society of Gene Therapy, Seattle, Washington.

CLAIMS

1. A gene delivery vehicle having been provided with at least a tissue tropism for smooth muscle cells and/or endothelial cells.
2. A gene delivery vehicle having been deprived of at least a tissue tropism for liver cells.
3. A vehicle according to claim 1 wherein said vehicle has been deprived of at least a tissue tropism for liver cells.
4. A vehicle according to anyone of the claims 1-3, wherein said tissue tropism is being provided by a virus capsid.
5. A vehicle according to claim 4, wherein said capsid comprises protein fragments from at least two different viruses.
6. A vehicle according to claim 5, wherein at least one of said viruses is an adenovirus.
7. A vehicle according to claim 5 or claim 6, wherein at least one of said viruses is an adenovirus of subgroup B.
8. A vehicle according to anyone of the claims 5-7, wherein at least one of said protein fragments comprises a tissue tropism determining fragment of a fiber protein derived from a subgroup B adenovirus.
9. A vehicle according to anyone of the claim 7 or claim 8, wherein said subgroup B adenovirus is adenovirus 16.
10. A vehicle according to claim 7-9, wherein protein fragments not derived from an adenovirus of subgroup B are derived from an adenovirus of subgroup C, preferably of adenovirus 5.
11. A vehicle according to anyone of the claims 1-10 comprising a nucleic acid derived from an adenovirus.
12. A vehicle according to anyone of the claims 1-11, comprising a nucleic acid derived from at least two different adenoviruses.

13. A vehicle according to claim 11 or claim 12, wherein said nucleic acid comprises at least one sequence encoding a fiber protein comprising at least a tissue tropism determining fragment of a subgroup B adenovirus fiber protein, preferably of adenovirus 16.
14. A vehicle according anyone of the claims 10-13, wherein said adenovirus nucleic acid is modified such that the capacity of said adenovirus nucleic acid to replicate in a target cell has been reduced or disabled.
15. A vehicle according to anyone of the claims 11-14, wherein said adenovirus nucleic acid is modified such that the capacity of a host immune system to mount an immune response against adenovirus proteins encoded by said adenovirus nucleic acid has been reduced or disabled.
16. A vehicle according to anyone of the claims 1-15, comprising a minimal adenovirus vector or an Ad/AAV chimaeric vector.
17. A vehicle according to anyone of the claims 1-16, further comprising at least one non-adenovirus nucleic acid.
18. A vehicle according to claim 17 wherein at least one of said non-adenovirus nucleic acids is a gene selected from the group of genes encoding: an apolipoprotein, a nitric oxide synthase, a herpes simplex virus thymidine kinase, an interleukin-3, an interleukin-1 $\alpha$ , an (anti)angiogenesis protein such as angiostatin, an anti-proliferation protein, a smooth muscle cell anti-migration protein, a vascular endothelial growth factor (VGEF), a basic fibroblast growth factor, a hypoxia inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ) or a PAI-1.
19. A cell for the production of a vector according to anyone of the claims 1-18, comprising means for the assembly of said vectors wherein said means includes a means for the production of an adenovirus fiber protein, wherein said fiber protein comprises at least a tissue tropism determining fragment of a subgroup B adenovirus fiber protein.
20. A cell according to claim 19, wherein said cell is or is derived from a PER.C6 cell (ECACC deposit number 96022940).

21. The use of a vehicle according to anyone of the claims 1-18 as a pharmaceutical.
22. The use of claim 21 for the treatment of cardiovascular disease.
- 5 23. The use of claim 21 for the treatment of a disease, treatable by transfer of a therapeutic nucleic acid to smooth muscle cells and/or endothelial cells.
24. An adenovirus capsid with or provided with a tissue tropism for smooth muscle cells and/or endothelial cells
- 10 wherein said capsid preferably comprises proteins from at least two different adenoviruses and wherein at least a tissue tropism determining fragment of a fiber protein is derived from a subgroup B adenovirus, preferably of adenovirus 16.
- 15 25. An adenovirus capsid having been deprived of a tissue tropism for liver cells wherein said capsid preferably comprises proteins from at least two different adenoviruses and wherein at least a tissue tropism determining fragment of a fiber protein is derived from a subgroup B adenovirus,
- 20 preferably of adenovirus 16.
26. The use of an adenovirus capsid according to claim 24 and/or claim 25, for the delivery of nucleic acid to smooth muscle cells and/or endothelial cells.
27. The use of an adenovirus capsid according to claim 26,
- 25 in a medicament for the treatment of a disease.
28. Construct pBr/Ad.BamRAFib, comprising adenovirus 5 sequences 21562-31094 and 32794-35938.
29. Construct pBr/AdBamRfib16, comprising adenovirus 5 sequences 21562-31094 and 32794-35938, further comprising an
- 30 adenovirus 16 gene encoding fiber protein.
30. Construct pBr/AdBamR.pac/fib16, comprising adenovirus 5 sequences 21562-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein, and further comprising a unique PacI-site in the proximity of the
- 35 adenovirus 5 right terminal repeat, in the non-adenovirus sequence backbone of said construct.

31. Construct pWE/Ad.AflIIrITRfib16, comprising adenovirus 5 sequences 3534-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein.
32. Construct pWE/Ad.AflIIrITRDE2Afib16, comprising  
5 adenovirus 5 sequences 3534-22443, 24033-31094 and 32794-35938, further comprising an adenovirus 16 gene encoding fiber protein.
33. The use of a construct according to anyone of the claims 28-32 for the generation of a vehicle according to  
10 anyone of the claims 1-18 or an adenovirus capsid according to claim 24 or claim 25.
34. The production of a vehicle according to anyone of the claims 1-18 or a adenovirus capsid according to claim 24 or claim 25.
- 15 35. The use of a vehicle according to anyone of the claims 1-18 for the generation a gene library.
36. The use of a fiber protein of adenovirus 16 for the delivery of nucleic acid to smooth muscle cells and/or endothelial cells.
- 20 37. The use of a fiber protein of adenovirus 16 in an adenovirus capsid for depriving said capsid of a tissue tropism for liver cells.

1/24





Figure 1

2/24

Figure 2



3/24




Figure 3

4/24

Figure 4A: Sequence of Ad5 fiber

ATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATA/TGACACGGAA/ACCGGTC  
CTCCAACGTGTCCTTTTCTTACTCCTCCCTTTGTATCCCCCAATGGGTTTCAAGAGAGTCCCCCTGG  
GGTACTCTCTTTGCGCCTATCCGAACCTCTAGTTACCTCCAATGGCATGCTTGCGCTCAAAATGGGC  
AACGGCCTCTCTCTGGACGAGGCCGGCAACCTTACCTCCCCAAATGTAACCACTGTGAGCCCACCTC  
TCAAAAAAACCAAGTCAAACATAAACCTGGAAATATCTGCACCCCTCACAGTTACCTCAGAAGCCCT  
AACTGTGGCTGCGCGCGCACCTCTAATGGTCGCGGGCAACACACTCACCATGCAATCACAGGCCCCG  
CTAACCGTGCACGACTCCAAACTTAGCATTGCCACCCAAGGACCCCTCACAGTGTGAGAAGGAAAGC  
TAGCCCTGCAAAACATCAGGCCCCCTCACCACCACCGATAGCAGTACCCTTACTATCACTGCCTCACC  
CCCTCTAACTACTGCCACTGGTAGCTTGGGCATTGACTTGAAAGAGCCCATTTATACACAAAATGGA  
AACTAGGACTAAAGTACGGGGCTCCTTTGCATGTAACAGACGACCTAAACACTTTGACCGTAGCAA  
CTGGTCCAGGTGTGACTATTAATAATACTTCCCTTGCAAACTAAAGTTACTGGAGCCTTGGGTTTGA  
TTCACAAGGCAATATGCAACTTAATGTAGCAGGAGGACTAAGGATTGATTCTCAAAACAGACGCCTT  
ATACTTGATGTTAGTTATCCGTTTGATGCTCAAACCAACTAAATCTAAGACTAGGACAGGGCCCTC  
TTTTTATAAACTCAGCCCACTTGGATATTAACCTACAACAAAGGCCTTTACTTGTGTACAGCTTC  
AAACAATTCCAAAAGCTTGAGGTTAACCTAAGCACTGCCAAGGGGTTGATGTTTGACGCTACAGCC  
ATAGCCATTAATGCAGGAGATGGGCTTGAATTTGGTTTACCTAATGCACCAAAACACAAATCCCCCTCA  
AAACAAAATTTGGCCATGGCCTAGAATTTGATTCAAACAAGGCTATGGTTCTAACTAGGAACTGG  
CCTTAGTTTTGACAGCACAGGTGCCATTACAGTAGGAAACAAAATAATGATAAGCTAACTTTGTGG  
ACCACACCAGCTCCATCTCCTAACTGTAGACTAAATGCAGAGAAAGATGCTAAACTCACTTTGGTCT  
TAACAAAATGTGGCAGTCAAATACTTGCTACAGTTTCAGTTTTGGCTGTTAAAGGCAGTTTGGCTCC  
AATATCTGGAACAGTTCAAAGTGCTCATCTTATTATAAGATTTGACGAAAATGGAGTGCTACTAAAC  
AATTCTCTTCTGGACCCGAATATTGGAACTTTAGAAATGGAGATCTTACTGAAGGCACAGCCTATA  
CAAACGCTGTTGGATTTATGCCTAACCTATCAGCTTATCCAAAATCTCACGGTAAACTGCCAAAAG  
TAACATTGTCACTCAAGTTTACTTAAACGGAGACAAAATAACCTGTAACACTAACCATTACACTA  
AACGGTACACAGGAAACAGGAGACACAACCTCCAAGTGCATACTCTATGTCATTTTCATGGGACTGGT  
CTGGCCACAACCTACATTAAATGAAATATTTGCCACATCCTCTTACACTTTTTCATACATTGCCCAAGA  
ATAA

Figure 4B: Sequence of Ad5/fib12 chimeric fiber

ATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATATGACCCATTTGACACAT  
CAGACGTACCCTTTGTTACACCCCTTTTACTTCTTCCAATGGTCTTCAAGAAAAACCACCGGTGT  
ATTAGCACTTAATTACAAAGACCCCATTTGTAAGTGAATGGAACCTTACACTCAAGCTAGGGGAC  
GGAATAAACTTAATGCCCAAGGTCAACTTACAGCTAGTAATAATATCAATGTTTTGGAGCCCTTA  
CCAACACCTCACAAGGTCTTAACTTTCTTGGAGCGCCCCCTAGCAGTAAAGGCTAGTGCCCTCAC  
ACTTAACACAAGAGCGCCCTTAACCACAACGGATGAAAGCTTAGCCTTAATAACCGCCCTCCCATTT  
ACAGTAGAGTCTTCGCGTTTGGGCTTGGCCACCATAGCCCTCTAAGCTTAGATGGAGGTGGAACC  
TAGGTTTAAATCTTTCTGCTCCCTGGACGTTAGTAACAACAATTTGCATCTCACCCTGAACTCC  
CTTAGTTGTAATCTAGCGGTGCCCTATCTGTTGCTACTGCAGACCCATAAGTGTTTCGCAACAAC  
GCTCTTACCCTACCTACGGCAGATCCGTTAATGGTGAGCTCCGATGGGTGGGAATAAGTGTCACATA  
GTCCCATTACAGTAATAAACGGTTCCCTTAGCCTTGTCTACAACCTGCTCCCTCAACAGCACAGGATC  
CACTTTAAGTCTGTCTGTTGCCAATCCTCTGACTATTTTACAAGACACATTGACTGTTTCCACTGGT  
AACGGTCTTCAAGTGTGGGGTCTCAATTAGTAACAAGAATAGGGGATGGTTTAACATTTCGATAATG  
GGGTCATGAAAGTAAACGTTGCCGGGGGAATGAGAAGTCTGGCGGTAGAATAATTTTAGATGTTAA  
TTATCCCTTTGATGCGAGCAATAACCTGTCTTAAGACGGGGATTGGGACTAATTTATAACCAATCT  
ACAACTGGAACCTTAACAACCTGATATTAGTACCGAAAAAGGTTTAAATGTTTAGTGGCAATCAATAG  
CTCTTAATGCAGGTCAGGGGCTTACATTTAATAATGGCCAACCTTAGGGTTAAGTTGGGAGCTGGACT  
TATTTTTGATTCAAACAATAACATTGCCTTAGGCAGCAGCAGCAACACTCCATACGACCTCTGACA  
CTGTGGACAACCTCTGACCCACCACCAAACTGCAGCCTCATACAAGAGCTAGATGCAAACTCACC  
TGTGCTTAACAAAAACGGATCTATTGTTAATGGCATTGTAAGTTTAGTGGGTGTTAAGGGTAATCT  
CCTAAATATCCAAAGTACTACTACCCTGTAGGAGTGCAATTTAGTGTTTGATGAACAGGGAAGATTA  
ATCACATCAACCCCTACTGCCCTGGTTCCCAAGCTTCGTGGGGATATAGACAAGGCCAATCAGTGT  
CTACCAATACTGTTACCAATGGTCTAGGTTTTATGCCTAATGTGAGTGCTTACCCTAGACCAATGC  
CAGTGAGGCTAAAAGCCAAATGGTAAGTCTCAGTACTTACAGGGAGATACATCTAAACCTATAACA  
ATGAAAGTTGCATTTAATGGCATTACGTCGCTAAATGGATACTCTTTAACATTCATGTGGTCAGGTC  
TATCAAACTATATAAATCAGCCTTTCTCTACACCATCCTGCTCCTTNTCTTACATTGCCCAAGAAATA  
AATGCATTAG

6/24

Figure 4C: Sequence of Ad5/fib16 chimeric fiber

ATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATATGAAGATGAAAGCAGCT  
CACAACACCCCTTTATAAACCCCTGGTTTCATTTCTCTCAAATGGTTTTGCACAAAGCCCAGATGGAGT  
TCTAACTCTTAAATGTGTTAATCCACTCACTACCGCCAGCGGACCCCTCCAACCTTAAAGTTGGAAGC  
AGTCTTACAGTAGATACTATCGATGGGTCTTTGGAGGAAAATATAACTGCCGAAGCGCCACTCACTA  
AAACTAACCCTCCATAGGTTTATTAATAGGATCTGGCTTGCAAACAAAGGATGATAAACTTTGTTT  
ATCGCTGGGAGATGGGTGGTAACAAAGGATGATAAACTATGTTTATCGCTGGGAGATGGGTAAATA  
ACAAAAATGATGTACTATGTGCCAACTAGGACATGGCCTTGTGTTTGACTCTTCCAATGCTATCA  
CCATAGAAAACAACACCTTGTGGACAGGCGCAAAACCAAGCGCCAACCTGTGTAATTAAAGAGGGAGA  
AGATTCCCCAGACTGTAAGCTCACTTTAGTTCTAGTGAAGAATGGAGGACTGATAAATGGATACATA  
ACATTAATGGGAGCCTCAGAATATACTAACACCTTGTTTAAAAACAATCAAGTTACAATCGATGTAA  
ACCTCGCATTTGATAATACTGGCCAAATTATTACTTACCTATCATCCCTTAAAGTAACCTGAACCTT  
TAAAGACAACCAAAACATGGCTACTGGAACCATAACCAGTGCCAAAGGCTTCATGCCCAGCACCACC  
GCCTATCCATTTATAACATACGCCACTGAGACCCTAAATGAAGATTACATTTATGGAGAGTGTTACT  
ACAAATCTACCAATGGAACCTCTTTCCACTAAAAGTTACTGTCACTAAACAGACGTATGTTAGC  
TTCTGGAATGGCCTATGCTATGAATTTTTCATGGTCTCTAAATGCAGAGGAAGCCCCGGAAACTACC  
GAAGTCACTCTCATTACCTCCCCCTCTTTTTTCTTATATCAGAGAAGATGACTGAATGCATTAG

7/24

Figure 4D: Sequence of Ad5/fib28 chimeric fiber

ATGTTGTTGCAGATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATATGGCT  
ACGCGCGGAATCAGAATATCCCCTTCCTCACTCCCCCTTTGTTTCTTCCGATGGATTCCAAAACCTT  
CCCACCTGGGGTCTGTCACTCAAACCTGGCTGACCCAATCACCATCGCTAATGGGGATGTCTCACTC  
AAGTTGGGAGGCGGACTGACGGTGGAAAAAGAGTCTGGAACTTAAGTGTGAACCCCTAAGGCTCCCT  
TGCAAGTTGCAAGTGGACAATTGGAATTAGCATATGATTCTCCATTTGATGTTAAAAACAATATGCT  
TACTCTTAAAGCAGGTCACGGCTTAGCAGTTGTAACGAAAGACAATACTGATTTACAACCACTAATG  
GGCACACTTGTGTTTTAACTGGCAAAGGCATTGGCACTGGCACAAGTGCTCACGGTGGAAACCATAG  
ATGTGAGAATAGGAAAAACGGAAGTCTGGCATTGACAAAAATGGAGATTTGGTGGCCTGGGATAA  
AGAAAATGACAGGCGCACTCTATGGACAACCTCAGACACATCTCCAAATTGCAAAATGAGTGAAGTC  
AAAGACTCAAAGCTTACTCTTATTCTTACAAAATGCGGAAGTCAAATTCTAGGAAGTGATCTTTGC  
TTGCTGTAAAAGGAGAATATCAAAATATGACTGCCAGTACTAATAAGAATGTAAAAATAACACTGCT  
ATTTGATGCTAATGGAGTCTTGTTAGAAGGATCCAGTCTTGATAAAGAGTACTGGAACCTTAGAAAC  
AATGATTCTACTGTGTCTGGAAAATATGAAAATGCTGTTCCGTTTCATGCCTAACATAACAGCTTATA  
AACCCGTCAATTCTAAAAGCTATGCCAGAAGTCACATATTTGGAAATGTATATATTGCTGCTAAGCC  
ATATAATCCAGTGGTTATTAAAATTAGCTTCAATCAAGAGACACAAAACAATTGTGTCTATTCTATA  
TCATTTGACTACACTTGCTCTAAAGAGTATACAGGTATGCAATTCGATGTTACATCTTTCACCTTCT  
CCTATATCGCCCAAGAATGAATGCATTAG

8/24

Figure 4E: Sequence of Ad5/fib40-L chimeric fiber

ATGTTGTTGCAGATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATATGAAC  
ACTACAATCCCCTTGACATTCATTTATTACACCCCGTTTGCTTCCTCCAACGGCTTGCAAGAAAA  
ACCTCCGGGAGTCCTCAGCCTGAAATACACTGATCCACTTACAACCAAAAACGGGGCTTTAACCTTA  
AAATTGGGACACGGGACTAAACATTGATAAAAAATGGAGATCTTTCTTCAGATGCTAGCGTGGAAGTTA  
GCGCCCTATCACTAAACCAACAAAATCGTAGGTTTAAATTACACTAAGCCTCTCGCTCTGCAAAA  
TAACGCGCTTACTCTTTCTTACAACGCGCCCTTTAACGTAGTAAATAATAATTTAGCTCTAAATATG  
TCACAGCCTGTTACTATTAATGCAACAACGAACCTTTCTCTCTTAATAGACGCCCCACTTAATGCTG  
ACACGGGCACTCTTCGCCCTTCGAAGTGATGCACCTCTTGGAAGTAGTAGACAAAACACTAAAGGTTT  
GTTTTCTAGCCCCCTCTATCTAGATAATAACTTTCTTACACTAGCCATTGAACGCCCCGCTAGCTCTA  
TCCAGTAACAGAGCAGTGCCCTTAAGTATTCACCACCTTTAAAAATAGAAAACGAAAACCTTAACCC  
TAAGCACAGGCGGACCTTTTACTGTAAAGCGGGGAAATTTAAACCTGGCAACATCGGCACCCCTCTC  
CGTGCAAAAACAATTCTCTCTCCTTAGGGGTTAACCCGCTTTTCTCATCACTGACTCTGGATTAGCT  
ATGGACTTAGGAGACGGTCTTGCAATTAGGTGGCTCTAAGTTAATAATCAATCTTGGTCCAGGTTTAC  
AAATGTCTAATGGAGCTATTACTTTAGCACTAGATGCAGCGCTGCCTTTGCAATATAAAAACAACCA  
ACTTCAACTCAGAATTGGCTCCGCGTCTGCTTTAATTATGAGCGGAGTAACACAAACATTAAACGTC  
AATGCCAATACCAGCAAAGGTCTTGCTATTGAAAATAACTCACTAGTTGTTAAGCTAGGAAACGGTC  
TTCGCTTTGATAGCTGGGGAAGCATAGCTGTCTCACCTACTACCACTACCCCTACCACCTATGGAC  
CACC CGGACCCGCTCTCCTAACGCCACTTTTTATGAATCACTAGACGCCAAAGTGTTGGCTAGTTTTA  
GTAAAATGCAACGGCATGGTTAACGGGACCATATCCATTAAAGCTCAAAAAGGCACCTTACTTTAAAC  
CCACAGCTAGCTTTATTTCTTTGTCTATGTTTTTACAGCGACGGAACGTGGAGGAAAACTATCC  
CGTGTGTTGACAACGAAGGGATACTAGCAAAACAGTGCCACATGGGGTTATCGACAAGGACAGTCTGCC  
AACACTAACGTTTCCAATGCTGTAGAATTTATGCCTAGCTCTAAAAGGTATCCCAATGAAAAAGGTT  
CTGAAGTTCAGAACATGGCTCTTACCTACACTTTTTTGAAGGTGACCCTAACATGGCCATATCTTT  
TCAGAGCATTATAATCATGCAATAGAAGGCTACTCATTAAAAATTCNCCTGGCGCGTTTCGAAATAAT  
GAACGTTTTGACATCCCCTGTTGCTCATTTTCTTATGTAAACAGAACAATAAATGCATTAG

9/24





Figure 5


10/24

Figure 6: Generation of (chimera) adenoviruses




11/24

Figure 7a




12/24

Figure 7b




13/24

Figure 7c




14/24

Figure 8a




15/24

Figure 8b




16/24

Figure 8c




17/24

Figure 8d



18/24


Figure 8e:



Ad5Fiber 51.ntLacZ



Ad5Fiber 16.ntLacZ




Ad5.ntLacZ



Negative control


19/24

Figure 8f



20/24

Figure 8g



Ad5Fiber 16.ntLacZ



Ad5.ntLacZ



Negative control

21/24

Figure 8h






Figure 9A

Alignment Report of Untitled, using Clustal method with Weighted residue weight table.  
Thursday, November 19, 1998 18:25

```
1  ATGGC---CAAACGAGCTCGGCTAAGCAGCT----- Ad16 genbank.se
1  ATGTTGTTGCA[GA]TGAAGCGCGC[AAGA]C[G]TCTGAAGATA Ad5/fib16.seq

29  CCTTCAATCCGGTCTACCCCTATGAAGATGAAAGCAGCTC Ad16 genbank.se
41  CCTTCAA[CC][CGT][GTA]TCC[A]TATGAAGATGAAAGCAGCTC Ad5/fib16.seq

69  ACAACACCCCTTTATAAACCCCTGGTTTTCATTTCTCTCAAAT Ad16 genbank.se
81  ACAACACCCCTTTATAAACCCCTGGTTTTCATTTCTCTCAAAT Ad5/fib16.seq

109 GGTTTTGCACAAAGCCCAGATGGAGTTCTAACTCTTAAAT Ad16 genbank.se
121 GGTTTTGCACAAAGCCCAGATGGAGTTCTAACTCTTAAAT Ad5/fib16.seq

149 GTGTTAATCCACTCACTACCGCCAGCGGACCCCTCCAAT Ad16 genbank.se
161 GTGTTAATCCACTCACTACCGCCAGCGGACCCCTCCAAT Ad5/fib16.seq

189 TAAAGTTGGAAGCAGTCTTACAGTAGATACTATCGATGGG Ad16 genbank.se
201 TAAAGTTGGAAGCAGTCTTACAGTAGATACTATCGATGGG Ad5/fib16.seq

229 TCTTTGGAGGAAAAATATAACTGCCGAGCGCCACTCACTA Ad16 genbank.se
241 TCTTTGGAGGAAAAATATAACTGCCG[A]AGCGCCACTCACTA Ad5/fib16.seq

269 AAATAAACCCTCCATAGGTTTATTAATAGGATCTGGCTT Ad16 genbank.se
281 AAATAAACCCTCCATAGGTTTATTAATAGGATCTGGCTT Ad5/fib16.seq

309 GCAAACAAAGGATGATAAACTTTGTTTATCGCTGGGAGAT Ad16 genbank.se
321 GCAAACAAAGGATGATAAACTTTGTTTATCGCTGGGAGAT Ad5/fib16.seq

349 GGGTTGGTAACAAAGGATGATAAACTATGTTTATCGCTGG Ad16 genbank.se
361 GGGTTGGTAACAAAGGATGATAAACTATGTTTATCGCTGG Ad5/fib16.seq

389 GAGATGGGTAAATAACAAAAAATGATGTACTATGTGCCAA Ad16 genbank.se
401 GAGATGGGTAAATAACAAAAAATGATGTACTATGTGCCAA Ad5/fib16.seq

429 ACTAGGACATGGCCCTTGTGTTTGA CTCTTCCAATGCTATC Ad16 genbank.se
441 ACTAGGACATGGCCCTTGTGTTTGA CTCTTCCAATGCTATC Ad5/fib16.seq

469 ACCATAGAAAAACAACACCTTGTGGACAGGCGCAAAACCAA Ad16 genbank.se
481 ACCATAGAAAAACAACACCTTGTGGACAGGCGCAAAACCAA Ad5/fib16.seq

509 GCGCCAACCTGTGTAATTAAAGAGGGAGAAGATTCCCCAGA Ad16 genbank.se
521 GCGCCAACCTGTGTAATTAAAGAGGGAGAAGATTCCCCAGA Ad5/fib16.seq

549 CTGTAAGCTCACTTTAGTTCTAGTGAAGAATGGAGGACTG Ad16 genbank.se
561 CTGTAAGCTCACTTTAGTTCTAGTGAAGAATGGAGGACTG Ad5/fib16.seq

589 ATAAATGGATACATAACATTAATGGGAGCCTCAGAATATA Ad16 genbank.se
601 ATAAATGGATACATAACATTAATGGGAGCCTCAGAATATA Ad5/fib16.seq

629 CTAACACCTTGTTTAAAAACAATCAAGTTACAATCGATGT Ad16 genbank.se
641 CTAACACCTTGTTTAAAAACAATCAAGTTACAATCGATGT Ad5/fib16.seq

669 AAACCTCGCATTTGATAATACTGGCCAAATTATTACTTAC Ad16 genbank.se
681 AAACCTCGCATTTGATAATACTGGCCAAATTATTACTTAC Ad5/fib16.seq

709 CTATCATCCCTTAAAAGTAACCTGAACTTTAAAGACAACC Ad16 genbank.se
721 CTATCATCCCTTAAAAGTAACCTGAACTTTAAAGACAACC Ad5/fib16.seq
```

23/24

Figure 9A , contd.

Alignment Report of Untitled, using Clustal method with Weighted residue weight table.  
Thursday, November 19, 1998 18:26

749	AAAACATGGCTACTGGAACCATAACCAGTGCCAAAGGCTT	Ad16 genbank.se
761	AAAACATGGCTACTGGAACCATAACCAGTGCCAAAGGCTT	Ad5/fib16.seq
789	CATGCCCAGCACCAACCGCCTATCCATTTATAACATACGCC	Ad16 genbank.se
801	CATGCCCAGCACCAACCGCCTATCCATTTATAACATACGCC	Ad5/fib16.seq
829	ACTGAGACCCTAAATGAAGATTACATTTATGGAGAGTGTT	Ad16 genbank.se
841	ACTGAGACCCTAAATGAAGATTACATTTATGGAGAGTGTT	Ad5/fib16.seq
869	ACTACAAATCTACCAATGGAACCTCTCTTTCCACTAAAAGT	Ad16 genbank.se
881	ACTACAAATCTACCAATGGAACCTCTCTTTCCACTAAAAGT	Ad5/fib16.seq
909	TACTGTCACACTAAACAGACGTATGTTAGCTTCTGGAATG	Ad16 genbank.se
921	TACTGTCACACTAAACAGACGTATGTTAGCTTCTGGAATG	Ad5/fib16.seq
949	GCCTATGCTATGAATTTTTCATGGTCTCTAAATGCAGAGG	Ad16 genbank.se
961	GCCTATGCTATGAATTTTTCATGGTCTCTAAATGCAGAGG	Ad5/fib16.seq
989	AAGCCCCGGAAACTACCGAAGTCACTCTCATTACCTCCCC	Ad16 genbank.se
1001	AAGCCCCGGAAACTACCGAAGTCACTCTCATTACCTCCCC	Ad5/fib16.seq
1029	CTTCTTTTTTTTCTTATATCAGAGAAGATGACTGA	Ad16 genbank.se
1041	CTTCTTTTTTTTCTTATATCAGAGAAGATGACTGA	Ad5/fib16.seq

Decoration 'Decoration #1': Box residues that differ from Ad16 genbank.seq.

24/24

Figure 9B

Alignment Report of Untitled, using Clustal method with PAM250 residue weight table.  
Thursday, November 19, 1998 18:09

```

1  M A K R A R L S S [ ] S F N P V Y P Y E D E S S Q H P F I N Ad16 fiber protein GenBank
1  M [ ] K R A R [P] S [E] D [T] F N P V Y P Y E D E S S Q H P F I N Ad16A fib protein

30 P G F I S S N G F A Q S P D G V L T L K C V N P L T T A S G Ad16 fiber protein GenBank
30 P G F I S S N G F A Q S P D G V L T L K C V N P L T T A S G Ad16A fib protein

60 P L Q L K V G S S L T V D T I D G S L E E N I T A A A P L T Ad16 fiber protein GenBank
60 P L Q L K V G S S L T V D T I D G S L E E N I T A [E] A P L T Ad16A fib protein

90 K T N H S I G L L I G S G L Q T K D D K L C L S L G D G L V Ad16 fiber protein GenBank
90 K T N H S I G L L I G S G L Q T K D D K L C L S L G D G L V Ad16A fib protein

120 T K D D K L C L S L G D G L I T K N D V L C A K L G H G L V Ad16 fiber protein GenBank
120 T K D D K L C L S L G D G L I T K N D V L C A K L G H G L V Ad16A fib protein

150 F D S S N A I T I E N N T L W T G A K P S A N C V I K E G E Ad16 fiber protein GenBank
150 F D S S N A I T I E N N T L W T G A K P S A N C V I K E G E Ad16A fib protein

180 D S P D C K L T L V L V K N G G L I N G Y I T L M G A S E Y Ad16 fiber protein GenBank
180 D S P D C K L T L V L V K N G G L I N G Y I T L M G A S E Y Ad16A fib protein

210 T N T L F K N N Q V T I D V N L A F D N T G Q I I T Y L S S Ad16 fiber protein GenBank
210 T N T L F K N N Q V T I D V N L A F D N T G Q I I T Y L S S Ad16A fib protein

240 L K S N L N F K D N Q N M A T G T I T S A K G F M P S T T A Ad16 fiber protein GenBank
240 L K S N L N F K D N Q N M A T G T I T S A K G F M P S T T A Ad16A fib protein

270 Y P F I T Y A T E T L N E D Y I Y G E C Y Y K S T N G T L F Ad16 fiber protein GenBank
270 Y P F I T Y A T E T L N E D Y I Y G E C Y Y K S T N G T L F Ad16A fib protein

300 P L K V T V T L N R R M L A S G M A Y A M N F S W S L N A E Ad16 fiber protein GenBank
300 P L K V T V T L N R R M L A S G M A Y A M N F S W S L N A E Ad16A fib protein

330 E A P E T T E V T L I T S P F F F S Y I R E D D [ ] Ad16 fiber protein GenBank
330 E A P E T T E V T L I T S P F F F S Y I R E D D [ ] Ad16A fib protein

```

Decoration 'Decoration #1': Box residues that differ from the Consensus.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/NL 99/00717

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/861 C12N15/62 C12N15/34 C12N5/10 C07K14/075  
A61K48/00 A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 20051 A (GENVEC INC ; WICKHAM THOMAS J (US); KOVESDI IMRE (US); BROUGH DOUGL) 5 June 1997 (1997-06-05)  * page 7, paragraph 2; page 35, paragraph 2; figure 9 *	1-4, 11, 17, 21-23, 34, 35
X	GALL J ET AL: "Adenovirus type 5 and 7 capsid chimera: fiber replacement alters receptor tropism without affecting primary immune neutralization epitopes" J VIROL., vol. 70, no. 4, April 1996 (1996-04), pages 2116-2123, XP002050655 cited in the application figure 7A  --- -/-	2, 4-8, 10-18, 21, 22, 24, 25, 27, 34, 35

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 April 2000

Date of mailing of the international search report

26.04.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Lonnoy, 0

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 99/00717

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 44121 A (MEHTALI MAJID ;LEGRAND VALERIE (FR); TRANSGENE SA (FR); BOULANGER) 8 October 1998 (1998-10-08)  * page 4, line 26 - page 5, line 11; page 26, line 1 - line 6 *	2,4-8, 10-18, 21,25, 27,34,35
X	WICKHAM T ET AL: "Targeted adenovirus gene transfer to endothelial and smooth muscle cells by using bispecific antibodies" J VIROL., vol. 70, no. 10, October 1996 (1996-10), pages 6831-6838, XP002911349 cited in the application	1,11,17, 34,35
Y	figures 2,7	3,4
X	WO 98 33929 A (KARAYAN LUCIE ;HONG SAW SEE (FR); BOULANGER PIERRE (FR); CENTRE NA) 6 August 1998 (1998-08-06)	2,11
Y	claims 15,23; figure 5	3,4
X	FALLAUX F ET AL: "New helper cells and matched early region 1-deleted adenovirus vectors prevent generation of replication-competent adenoviruses" HUM GENE THER., vol. 9, 1 September 1998 (1998-09-01), pages 1909-1917, XP002111070 cited in the application table 2	19,20
A	ROELVINK P ET AL: "The coxsackievirus-adenovirus receptor protein can function as a cellular attachment protein for adenovirus serotypes from subgroups A, C, D, E, and F" J VIROL., vol. 72, no. 10, October 1998 (1998-10), XP002111071 cited in the application	
A	WO 98 22609 A (ARMENTANO DONNA E ;GREGORY RICHARD J (US); GENZYME CORP (US); SMIT) 28 May 1998 (1998-05-28) cited in the application	

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/NL 99/ 00717

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 21-23, 26, 27, 36 and 37, as far as in vivo method are concerned, are directed to methods of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/NL 99/00717

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9720051 A	05-06-1997	US 5962311 A	05-10-1999
		US 5846782 A	08-12-1998
		US 5965541 A	12-10-1999
		AU 1086897 A	19-06-1997
		AU 4155597 A	06-03-1998
		BG 102554 A	30-04-1999
		BR 9612685 A	20-07-1999
		CZ 9801623 A	14-10-1998
		EP 0863987 A	16-09-1998
		EP 0920524 A	09-06-1999
		JP 11501219 T	02-02-1999
		NO 982417 A	06-07-1998
		PL 327038 A	09-11-1998
		SK 72298 A	02-12-1998
		WO 9807877 A	26-02-1998
		AU 4080497 A	06-03-1998
		EP 0920514 A	09-06-1999
		WO 9807865 A	26-02-1998
WO 9844121 A	08-10-1998	FR 2761688 A	09-10-1998
		FR 2761689 A	09-10-1998
		AU 7054798 A	22-10-1998
		EP 0991763 A	12-04-2000
WO 9833929 A	06-08-1998	FR 2758821 A	31-07-1998
		FR 2758822 A	31-07-1998
		AU 6104698 A	25-08-1998
		CA 2251036 A	06-08-1998
		EP 0900280 A	10-03-1999
WO 9822609 A	28-05-1998	US 5877011 A	02-03-1999
		AU 5455298 A	10-06-1998
		EP 0946742 A	06-10-1999

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**